

First report of Fusarium solani species complex as a causal agent of Erythrina variegata decline and death after gall formation by Quadrastichus erythrinae on Okinawa Island, Japan

Keiko Kuroda, Izumi Chuma, Takeo Kihara, Tsubasa Murakami, Kuya Takashina, Daiki Hiraoka & Norikazu Kameyama

Journal of General Plant Pathology

ISSN 1345-2630

Volume 83

Number 6


J Gen Plant Pathol (2017) 83:344-357

DOI 10.1007/s10327-017-0738-3



Your article is published under the Creative Commons Attribution license which allows users to read, copy, distribute and make derivative works, as long as the author of the original work is cited. You may self-archive this article on your own website, an institutional repository or funder's repository and make it publicly available immediately.

First report of *Fusarium solani* species complex as a causal agent of *Erythrina variegata* decline and death after gall formation by *Quadrastichus erythrinae* on Okinawa Island, Japan

Keiko Kuroda¹  · Izumi Chuma¹ · Takeo Kihara¹ · Tsubasa Murakami¹ · Kuya Takashina¹ · Daiki Hiraoka¹ · Norikazu Kameyama²

Received: 27 December 2016 / Accepted: 18 August 2017 / Published online: 21 September 2017
© The Author(s) 2017. This article is an open access publication

Abstract The defoliation and mortality of *Erythrina* spp. are increasing on the southern islands of Japan as well as in Taiwan. Although infestation by the gall wasp *Quadrastichus erythrinae* has been accepted as the cause of *Erythrina* decline, its role in the death of hosts has never been investigated. We thus studied the cause of the decline, focusing on physiological changes and any contributions by microorganisms. From declining and defoliated trees, *Fusarium* sp. within *F. solani* species complex (FSSC) clade 3 was primarily detected, especially from discolored xylem and necrotic phloem that had an odor. This fungus belongs to the Ambrosia *Fusarium* clade, a group of symbionts of ambrosia beetles that includes the causal agent of *Fusarium* dieback on avocado. No specific fungi were detected from twigs and leaves with *Q. erythrinae* galls. According to histological observations, the internal symptoms of declining *E. variegata* are similar to those of sudden death syndrome in soybeans, which is caused by *F. virguliforme*, a member of FSSC clade 2. One of the isolates of the *Fusarium* sp. isolated from *E. variegata* induced wilt and necrosis of cortex and phloem in the seedlings after artificial inoculation, and the inoculated strain was reisolated. The present results demonstrated that the isolate was pathogenic to *E. variegata* and may be a causal agent of *Erythrina* decline.

Keywords *Fusarium solani* species complex · *Erythrina variegata* · Sudden death · *Quadrastichus erythrinae* · Gall wasp · Okinawa · Japan

Introduction

The decline of *Erythrina* trees—abnormal defoliation, absence of flowering, and death—has been increasing in Japan and Taiwan (Uechi et al. 2007). Gall formation on leaves and petioles caused by the gall wasp *Quadrastichus erythrinae* Kim has been accepted as the cause of this syndrome and death (Rubinoff et al. 2010; Uechi et al. 2007). Infestation by this gall wasp, which is hypothesized to be native to Africa, began simultaneously during the 2000s in various areas of the world. The first report was from Taiwan in 2003 (Yang et al. 2004). In Japan, infestation was found in 2005 on an Indian coral tree, *Erythrina variegata* L., in Ishigaki Island, Okinawa Prefecture (Uechi et al. 2007). It was also found in Singapore (Kim et al. 2004), India (Faizal et al. 2006), Hawaii (Gramling 2005) and Florida in the United States (Reimer 2007), Vietnam (Uechi et al. 2007), China, the Philippines (Heu et al. 2008), and Brazil (Culik et al. 2014). To date, eight species and one subspecies among 110 *Erythrina* spp. have been reported as hosts.

Erythrina variegata, called “deigo” in Japanese, was introduced before the nineteenth century to the southern islands of Japan from India or Malaysia and has been commonly planted in parks and along the seashore and roadsides. Its wood is also used for traditional Ryukyu lacquerware. Injecting tree trunks with an insecticide (thiamethoxam, Izutsuya Chemical Industry) effectively kills larvae in the galls of *Q. erythrinae* (Kiyuna 2008); however, this technique did not decrease the mortality of *E. variegata*. Another species, *E. crista-galli* L., which is widely planted in the Honshu and Kyushu regions

✉ Keiko Kuroda
kurodak@garnet.kobe-u.ac.jp

✉ Izumi Chuma
chuizm@kobe-u.ac.jp

¹ Graduate School of Agricultural Science, Kobe University, 1-1 Rokkodai-cho, Nada-ku, Kobe City 657-8501, Japan

² Faculty of Agriculture, University of the Ryukyus, 1 Senbaru, Nishihara-cho, Nakagami-gun, Okinawa 903-0213, Japan

of Japan, is also under scrutiny. Due to these circumstances, a new strategy to protect *Erythrina* spp. is urgently needed.

Whether *Q. erythrinae* can directly kill trees is in question. In general, deciduous trees recover from unexpected defoliation by insects and seldom die from insect infestations. First, defoliation and the sudden death of host trees are disadvantageous to the propagation of *Q. erythrinae*. The sudden necrosis of bark tissue accompanied by an odor and the exfoliation of host trunks before death, when branches and twigs are still alive, suggest the activity of microorganisms in the lower part of the tree rather than trouble in the leaves. From a pathological perspective, *Q. erythrinae* may thus not be killing *E. variegata*. We obtained information that *Erythrina* death had occurred in Okinawa before 2005 without *Q. erythrinae* infestation, and the dead tree was processed as “cause unknown” (Y. Sakai, Okinawa Prefectural Government, personal communication). Unfortunately, there were no investigations on the relationship between *Q. erythrinae* infestation and symptom development in host trees. Previous reports on the mortality or wilt diseases of trees, such as Japanese oak wilt caused by *Raffaelea quercivora* (Kuroda 2001), fig wilt (fig canker) caused by *Ceratocystis ficicola* (Morita et al. 2016), and spruce wilt caused by *C. polonica* (Kuroda 2005), suggest that the contribution of microorganisms to *Erythrina* decline should also be checked. In preliminary investigations, *Fusarium* spp. were frequently isolated from the stems of declining but still living trees (Kihara et al. 2016). To develop techniques to protect *Erythrina* spp., we need to determine the specific cause of the decline and mortality of the hosts.

In the present investigation, we studied the physiology and pathology of living trees with defoliation to identify anomalies and determine the causal factors of *E. variegata* decline and mortality. The contribution of microbes to host mortality, physiological changes and internal symptoms such as the defense reaction of host cells (Hillis 1987), xylem dysfunction (Kuroda 2001; Morita et al. 2016), and the distribution of candidate microbes in the declining trees were also examined. Trees were then inoculated with candidate pathogens to confirm pathogenicity. In the case of large trees, an investigation of internal symptoms is very important because sometimes a time lag from infection to visible wilt conceals the real causal factor(s). This report includes preliminary information for quickly controlling *Erythrina* decline.

Materials and methods

Sampling and macroscopic observations

Erythrina variegata was sampled three times during 2014 and 2015 from trees 30 to 60 cm in diameter (Table 1). Heavily damaged trees were cut and harvested from the

base, or the largest branches were cut from planted trees that were not permitted to be cut down from the base. Two trees were harvested on May 20, 2014, at the University of the Ryukyus, Nisihara-cho, Nakagami-gun, Okinawa Prefecture, Japan. Specimen R1 had sparse leaves and was judged to be close to death (Level 1, extensive damage). A harvested, thick branch was cut into three pieces and used for microscopy and the isolation of microorganisms. Specimen R2 retained more leaves than on R1, although it was abnormally defoliated (Level 2, moderate damage). On September 9, 2014, one tree, P1, with few leaves (Level 1) was harvested in the Okinawa Prefectural Peace Memorial Park, Itoman City, Okinawa Prefecture. A thick branch 3.7 m long was cut into five pieces (Table 2). In addition, four branches with abundant leaves (Level 2), R3–R6, were harvested at the University of the Ryukyus. Galls of *Q. erythrinae* were not found on specimen R3 but were found on the petioles of R4 and R5 and on the leaf surface of R6. On July 29, 2015, specimen P2, with few leaves (Level 1) but without bark exfoliation, and specimen P3, with abundant leaves (Level 2), were selected at Peace Memorial Park. After the dye injection experiment explained below, the main trunk of P2 and a thick branch of P3 were harvested.

Healthy trees without infestation of *Q. erythrinae* do not exist in the natural environment in Okinawa Prefecture. Therefore, seedlings grown in a greenhouse without infestation of *Q. erythrinae* were provided from Okinawa Prefectural Forest Resource Research Center as control specimens. Portions of the 10 seedlings provided on December 1, 2014, were used for microscopy and for isolating microorganisms (Table 2 C1–C3).

Water conduction in xylem

To check xylem sap ascent, a dye solution was injected into the trunks of declining but living trees P2 (Level 1) and P3 (Level 2) on July 29, 2015, at Peace Memorial Park. Two holes, each 5 mm in diameter and 3 cm deep, were drilled in each trunk. A silicone tube was inserted into each hole, and 1% (w/v) aqueous acid fuchsin in a 1-L bottle was allowed to drip into the tree from 11 a.m. to 2 p.m. On a sunny day, dye solution will be automatically absorbed and water-conducting xylem dyed red in healthy trees. This technique is convenient for detecting the activity of xylem conduction visually and from the amount of absorbed liquid (Kuroda and Yamada 1996). The trunk and branch were cut from the base into 30- to 50-cm-long pieces so that stained areas on the cut surfaces could be observed by eye. Then the specimens were used for histological and pathological experiments.

Table 1 Sampling and experiments with declining *Erythrina variegata* trees

Sampling date	Tree no.	Sampling site	Tree height (m)	Diameter at breast height (cm)	Diameter of specimen (cm)	Leaf quantity level*	Condition, sampled part	Experimental purpose
May 20, 2014	R1	Univ. Ryukyus Nisihara-cho, Nakagami-gun	7.9	45.2	20.0	1	Heavy damage, thick branch	Histology Fungal isolation
	R2		8.7	44.8	8.0	2	Slight damage, thick branch 26-years-old	
Sept. 9, 2014	P1	Okinawa Prefectural Peace Memorial Park, Itoman City	7.8	58.6	29.0	1	Heavy damage, thick branch	Histology Fungal isolation
	R3		10.0	59.2	7.0	2	No galls, upper branch	
	R4		5.7	31.1	5.0	2	Galls on petioles, upper branch	
	R5		5.3	32.9	3.0	2	Galls on petioles, upper branch	
	R6		–	–	1.5	2	Galls on leaves, upper branch	
July 29, 2015	P2	Peace Memorial Park	5.2	48.1	–	1	Heavy damage, hole tree	Water conduction Histology Fungal isolation
	P3		6.4	48.4	–	2	Slight damage, thick branch	
Dec. 1, 2014	C1	Okinawa Pref. Forest Resource Research Center	0.20	–	0.5	3	No damage, potted seedlings 1 year old	Histology Fungal isolation
	C2		0.25	–	0.5	3		
	C3		0.24	–	0.5	3		

Level 1 small quantity of leaves, Level 2 abundant leaves, Level 3 healthy condition

Macro- and microscopic observations

Specimens were dissected into smaller pieces, and the cross-section and radial surfaces were observed with a stereoscopic microscope (SMZ1500, Nikon, Tokyo, Japan) and photographed with digital cameras. Specifically, the phloem and xylem of slightly damaged trees (Level 2) were compared with those of severely damaged trees (Level 1).

Next, specimens were processed for light microscopy. Some of the specimens were cut into small blocks (1 × 1 × 2 to 2 × 2 × 3 cm) that contained lesions or discoloration; these were fixed in FAA (formalin, acetic acid, 50% ethyl alcohol; 5:5:90, v/v) for 1 week and, subsequently, washed for 1 day under tap water. With a sliding microtome (Ritratome REM-710, Yamato Koki Co., Asaka, Saitama, Japan), 20- to 30-µm-thick sections (transverse, tangential, and radial) were cut from the small blocks. Parts of the sections were mounted onto slides without staining to observe the natural colors and cytological changes. Some sections were stained with safranin–fast green for the observation of lignified xylem cells and cell contents (Ruzin 1999). For the observation of fungal hyphae, parts of the sections were stained

with periodic acid–Schiff (PAS) and toluidine blue O (Feder and O'Brien 1968). Microscopic observation (ECLIPSE 80i, ECLIPSE Ni-U, Nikon) was focused on the range of discolored xylem, the distribution of fungal hyphae, and the necrosis of parenchyma cells.

Fungal isolation and identification

Specimens were cut from various parts of declining trees. In addition to abnormally discolored and necrotic phloem tissues, nondiscolored xylem and living phloem were examined for the presence of fungi (Table 2). In large diseased trees, the fungus is sometimes restricted to one part of the tree body. For instance, specimen R1 at the base of a branch contained discolored and nondiscolored xylem at the same height. Tissues were dissected into 1 × 3 × 5 mm cubes with disposable knife blades. Tissue cubes were washed with 70% ethanol and the surface was sterilized with aqueous sodium hypochlorite solution (effective chlorine concentration 0.5%), placed on potato dextrose agar (PDA, Difco) in 9-cm petri dishes, and incubated at 25 °C. In addition,

Table 2 Tissue condition in declining *Erythrina variegata* trees and healthy or inoculated seedlings, and fungi detected

Tree no.	Part examined	Xylem discolored ^a	Phloem necrosis ^a	Cultured part	No. of pieces cultured	<i>Fusarium</i> isolate tested		Other fungi isolated ^d
						Isolate ^b	Species ^c	
R1	Top of branch	–	–	Nondiscolored xylem Living phloem	42	R1-TB3	+	<i>Alternaria</i> sp. ^e <i>Papularia</i> sp. ^e
	Middle of branch	+	+	Discolored xylem	14	R1-MA7	+	<i>Papularia</i> sp. ^e
	Base of branch	+	+	Discolored xylem	63	R1-BC6	FSSC Strain B ^f	+
Nondiscolored xylem				21	–	–	<i>Papularia</i> sp. ^e	
R2	Branch	–	–	Nondiscolored xylem	42	–	–	<i>Papularia</i> sp. ^e <i>Pestalotia</i> sp. ^e
P1	Top of branch: 195 cm from base	–	+	Nondiscolored xylem	63	P1-160OA7	FSSC Strain A ^g	–
				Necrotic phloem	21	P1-195PA1	+	–
	90 cm from base	+	+	Discolored xylem	21	P1-90IB1	+	–
				Nondiscolored xylem	21	P1-90OA1	+	–
				Necrotic phloem	21	P1-90PA7	+	<i>Papularia</i> sp. ^e
	30 cm from base	+	+	Discolored xylem	21	–	–	<i>Papularia</i> sp. ^e
				Nondiscolored xylem	21	P1-30OA1	+	+
			Necrotic phloem	21	P1-30PA1	+	–	
R3	Upper branch	–	–	Nondiscolored xylem	42	R3-2XA1	+	+
				Living phloem				
R4	Upper branch	–	–	Petiole	10	R3-1LA1	+	+
				Nondiscolored xylem	42	R4-1PC6	+	<i>Nigrospora</i> sp. ^e , +
R5	Upper branch	–	–	Living phloem				
				Nondiscolored xylem	21	R5-1MA1	+	–
				Living phloem				
R6	Upper branch	–	–	Petiole	10	R5-2LA5	+	–
				Nondiscolored xylem	42	R6-SA1	+	–
P2 ^h	a: Branch	+	+	Nondiscolored xylem	21	–	–	<i>Colletotrichum</i> sp. ^g
				Living petiole	21	–	–	
	b: Branch	+	+	Discolored xylem	6	P2-a11	+	–
	c: Branch	+	+	Discolored xylem	6	P2-b11	+	<i>Lasiodiplodia</i> sp. ^g
	d: Branch	+	+	Discolored xylem	6	P2-c22	FSSC Strain A ^g	–
	e: Base of branch	+	–	Discolored xylem	6	P2-d11	FSSC Strain A ^g	<i>Lasiodiplodia</i> sp. ^g
	f: Lower trunk	+	+	Whitish hyphae on cross-cut surface	2	P2-e12	FSSC Strain A ^g	–
			Discolored xylem	8	P2-f22 (LC198904)	FSSC Strain B ^f	+	
					P2-f11 (LC198903, LC198905, LC198906, LC198907), P2-fK	FSSC Strain A ^g	+	
	g: Base of trunk	+	+	Discolored xylem	6	P2-g21	FSSC Strain B ^f	–

Table 2 (continued)

Tree no.	Part examined	Xylem discolored ^a	Phloem necrosis ^a	Cultured part	No. of pieces cultured	<i>Fusarium</i> isolate tested		Other fungi isolated ^d
						Isolate ^b	Species ^c	
P3 ^h	h: Top of branch	–	–	Nondiscolored xylem, living phloem	6	P3-h11	+	<i>Lasiodiplodia</i> sp. ^e
	i: Base of branch	+	–	Discolored xylem, living phloem	6	P3-i11	FSSC Strain B ^f	–
C1	Stem	–	–	Nondiscolored xylem, living phloem	21	–	–	–
	Petiole	–	–	Petiole	21	–	–	–
C2	Stem	–	–	Nondiscolored xylem, living phloem	21	–	–	–
	Petiole	–	–	Petiole	21	–	–	<i>Plectosphaerella</i> sp. ^g
C3	Stem	–	–	Nondiscolored xylem Living phloem	21	–	–	–
	Petiole	–	–	Petiole	21	–	–	+
AB2 ⁱ	Stem	–	+	Whitish hyphae on bark surface	2	AB2-S1	FSSC Strain A ^g	–
				Xylem, phloem	9	AB2-L3	FSSC Strain B ^f	–
A2 ⁱ	Root	–	–	Xylem, phloem	3	AB2-R1	FSSC Strain B ^f	–
	Stem	–	–	Xylem, phloem	9	–	–	<i>Myrothecium</i> sp. ^g
Ct2 ⁱ	Root	–	–	Xylem, phloem	3	A2-R2	FSSC Strain A ^g	–
	Stem	–	–	Xylem, phloem	9	A2-R3	FSSC Strain B ^f	–
Ct2 ⁱ	Stem	–	–	Xylem, phloem	9	–	+	+
	Root	–	–	Xylem, phloem	3	–	+	+

^aArea of xylem discoloration or phloem necrosis. +, detected; –, none

^bEMBL/GenBank accessions for nucleotide sequences are in parentheses

^cFSSC, *Fusarium solani* species complex; +, *Fusarium* sp. identified by conidial morphology; –, no fungus detected

^d+, Fungus isolated, genus not identified; –, no fungus detected

^eGenera identified by conidial morphology

^fStrain B, ITS nucleotide sequence was 99.8% (547/548 nt) identical to strain A

^gGenera identified by conidial morphology and ITS sequence

^hLetter codes a–i correspond to those in Fig. 1

ⁱInoculated seedlings. AB2, inoculated with P2-f11 and P2-f22; A2, inoculated with P2-f11; Ct2, no inoculation

hyphae that formed on the cut surface of some specimens within 5 days of the harvest date were picked with a lancet and incubated on PDA as above. Petioles of *E. variegata* obtained from trees R3, R5, and R6 were also sterilized and incubated as trunk tissue sections for detecting the site of infection. Control specimens cut from seedlings without *Q. erythrinae* infestation were also incubated on PDA. Spores formed on PDA were identified with a light microscope (Barnett and Hunter 1998). Strains that were predominantly isolated from declining trees were judged to be potential pathogens.

To identify species of the predominant isolates (Table 2), DNA sequencing of the nu-rRNA gene repeat (rDNA-ITS),

a portion of RNA polymerase II (*RPB2*), encoding the second largest subunit of RNA polymerase, and the translation elongation factor (*EF-1a*) gene was conducted as follows. Mycelial cultures grown on PDA were suspended in 100 µL of Tris–EDTA (TE) buffer, incubated at 95 °C for 10 min, and used as a PCR template. The rDNA-ITS was amplified with the primers ITS5 (5′-GGAAGTAAAAGT CGTAACAAGG-3′) and ITS4 (5′-TCCTCCGCTTATTGA TATGC-3′) (White et al. 1990), *RPB2* was amplified with primers 7cF (5′-ATGGG YAARCAAGCYATGGG-3′) and 11aR (5′-GCRTGGATCTTRTCRTCSACC-3′) (O’Donnell et al. 2007) and primers 5F2 (5′-GGGGWGAYCAGAAG AAGGC-3′) and 7cR (5′-CCCATRGC TTGYTTTRCCCAT-3′)

(O'Donnell et al. 2007), and *EF-1a* was amplified with primers EF-1H (5'-ATGGGTAAGGAAGACAAGAC-3') and EF-2T (5'-GGAAGTACCAGTGATCATGTT-3') (O'Donnell 2000) in a 20- μ L reaction of KOD FX Neo DNA polymerase (TOYOBO, Osaka, Japan). Sequence data for strains A and B are available in the EMBL/GenBank databases, with accession numbers listed in Table 2. We used the sequence data of *F. staphyleae* and 30 FSSC strains reported by Costa et al. (2016), Kasson et al. (2013), O'Donnell et al. (2008, 2015) and Zhang et al. (2006), which were obtained from GenBank for multiple sequence alignment using CLUSTAL W and analysis by MEGA7 (Kumar et al. 2016). The phylogenetic relationship was inferred by using the maximum likelihood method based on the Jukes–Cantor model (Jukes and Cantor 1969). All positions containing gaps and missing data were eliminated. Nodal supports were assessed using 1000 bootstrap replicates. The tree was rooted using *F. staphyleae* as an outgroup.

Inoculation experiments

Three inoculation experiments, including the preliminary one were done. Three seedlings were used for the preliminary experiment, and 8 and 11 seedlings were used for the main experiments, conducted twice (Table 3). Although inoculation experiments for tree diseases usually use 20–50 seedlings (Morita et al. 2016), fewer seedlings were used in the present experiments because seedlings of *E. variegata* are difficult to prepare since flowering and seed formation are rare in the Ryuku Islands. Twenty-two seedlings were provided by the Okinawa Prefectural Forest Resource Research Center on August 5, 2015, and 20 seedlings were provided by Ryukyu Sankei Co., Ltd., on June 2, 2016.

Inoculation schedules and specimen conditions are detailed in Table 3. In the preliminary experiment, a mycelial disk of three strains cultured for 2 weeks on PDA was placed onto a wound made at the base of seedlings (height ca. 30 cm) or petioles to narrow the number of candidates. Seedlings were observed for symptoms for 2 months. Two strains were chosen for the second inoculation experiment in 2015, and one of them was used for the third experiment in 2016.

The second experiment with 8 seedlings (height ca. 30 cm) was conducted on October 2, 2015. Soil was washed from the roots of seedlings under tap water. Then the seedlings were cultured in 200 mL of distilled water after their roots were cut to lengths of 6 cm. As inocula, spores of strain A (P2-f11) and strain B (P2-f22) were incubated on the PDA. Spores (1×10^5 /mL) of strains A, B, or A + B were added to two pots of water-cultured seedlings. Two more seedlings were cultured without fungal spores as controls. Symptom development was observed

for 2 months in an incubator set at 25 °C. Tissue of dead seedlings was used for reisolation.

In the third experiment, 11 seedlings were inoculated with strain A (P2-f11) on July 20, 2016 (Table 3). Mycelia incubated on sterilized toothpicks with PDA were placed in four small holes at the base of the lower stems of *E. variegata* seedlings.

Symptom development was checked every day or at 2-day intervals for 7–10 weeks. Death of specimens was judged as complete defoliation with no new leaves forming at the top or on the stem of seedlings. Tissue of dead seedlings was used for reisolations.

Results

Macroscopic characteristics of declining trees

Heavily damaged and almost defoliated *E. variegata* trees were commonly found adjacent to dead trees at both harvest sites at the University of the Ryukyus and Peace Memorial Park. Information from the local government of Okinawa Prefecture stated that the decline of trees was progressing outward from the center of a group of dead trees in Peace Memorial Park (Y. Sakai, Okinawa Prefectural Government, personal communication).

On the cut surface of the trunks and branches of heavily damaged and mostly defoliated specimens—R1, P1, and P2 (Level 1, Table 1; Fig. 1a)—pale brown or gray discolored xylem was visible (Fig. 2a, Dis: inside the dotted line). The discolored area was wider in the lower trunk of P2 (Fig. 2a, 82% of cross section) and in the lower part of a thick branch of R1. The bark of these trees, R1 and P1, was soft at various parts of the trunks and oozed liquid when pressed with fingertips. Necrotic phloem tissue in the softened area (Fig. 2, Nec) formed adjacent to the grayish discoloration in the xylem (Fig. 2a, Dis). A putrid smell was detected from the necrotic, softened tissue, and the necrotic bark tissue had exfoliated from the trunk surface. In trees with abundant leaves (Level 2, Table 1; Fig. 2b), branches of R2–R6, and the trunk of P3, discoloration and necrosis were not observed in the trunk tissue (Table 2).

Histological anomalies of declining trees

Changes in host cells and distribution of fungal hyphae

In the discolored xylem of heavily damaged trees—R1, P1, and P2 (Figs. 2a, 3a, 4a)—yellow to brown substances had accumulated in parenchyma cells and in the lumen of vessels (Fig. 3c), and the cell wall of the wood fibers

Table 3 Inoculation experiments with healthy seedlings of *Erythrina variegata*

Inoculation and harvest date (duration)	Specimen no.	Source of seedlings	Height (cm)	Diameter (cm)	Inoculation site
Preliminary July 21 to Sept. 10, 2015 (58 days)	L1	Okinawa Pref. Forest Resource Research Center	26–41	0.6–0.8	Petiole
	L2				Petiole
	L3				Petiole
	S1				Stem
	S2				Stem
	S3				Stem
	S4				Stem
	Oct. 2 to Nov. 17, 2015 (46 days)				AB1 ^a
AB2					
A1					
A2					
B1					
B2					
Ct1					
Ct2					
July 20 to Sept 28, 2016 (70 days)	1	Ryukyu Sankei Co., Ltd	21–54	0.7–1.5	Lower stem
	2				
	3				
	4				
	5				
	6				
	7				
	8				
	9				
	Ct1				
	Ct2				

^aSpecimen number includes the strain name. AB, A, or B means that inocula is a mixture of Strain A and B, or each one singly




Fig. 1 Condition of declining *Erythrina variegata* trees harvested at Peace Memorial Park in Itoman City, Okinawa Prefecture, on July 29, 2015. **a** Heavily damaged tree (P2). **b** Slightly damaged tree (P3). Arrows a–i sampling for histology and fungal isolation

was also stained. Fungal hyphae were observed in the discolored xylem area, vessels, axial parenchyma, wood fibers, and ray parenchyma tissues (Fig. 3b). Hyphae were abundant on the xylem side of the cambial region where the xylem discoloration had expanded and reached the cambium (Fig. 2a, Nec). Pale yellow substances were observed in some vessels and axial parenchyma of heavily damaged specimen P2 where slight discoloration was detected or indistinct by eye (Fig. 1a: c–e); hyphae were rarely found in such areas. In the cortex, mucilage ducts were filled with exudates.

In slightly deteriorated trees (abundant leaves, Level 2), R2–R5 and P3, no colored substances or hyphae were observed in the xylem. However, tyloses and oily substances were apparent in vessels and likely causing at least partial dysfunction of the vessels.

Leaves and petioles with galls of Q. erythrinae

When the histology of petioles with galls (R5) were compared with petioles without galls (R3) (Fig. 5), the




Fig. 2 Discoloration and dysfunction of xylem in the trunks of declining *E. variegata* trees harvested on July 29, 2015. **a** Wide xylem discoloration (Dis: inside the dotted line), necrosis of phloem and cambium (Nec), and extensive decrease of sap ascent (Sa) in the

heavily damaged tree (Fig. 1a, specimen P2). **b** Moderate sap ascent (arrows) in the slightly damaged tree (Fig. 1b, specimen P3). Water conduction was tested by the absorption of acid fuchsin solution for 3 h before the tree was cut down




Fig. 3 Colored substances and hyphae in discolored xylem of declining *E. variegata* in cross sections observed with light microscope. **a** Cross-cut surface of discolored xylem (Dis) in branch. **b** Hyphae (arrows) in ray tissue (R) and axial parenchyma cells (AP) of discolored area. **c** Pale-brown secondary metabolites, synthesized as a defense reaction to infection and filled in vessels (arrows). Section **b** was stained with toluidine blue O; section **c** was not stained

boundaries of the xylem, phloem, and cortex were distinct in the petioles of R3 (Fig. 5b). In contrast, in R5 tissue, numerous parenchymatous cells were abnormal and in disorderly arrangements, and the boundaries of the cortex and other

tissues were not distinguishable (Fig. 5c). No fungal hyphae or discolored host cells was observed around the galls.


Distribution and identity of fungi

In trunks and branches

Fusarium-type conidia with white to pale yellow mycelia (Table 2; Fig. 4) were identified from the predominant fungi isolated from various parts of declining trees. Other fungi such as the *Papularia* sp., *Pestalotia* sp., and *Alternaria* sp. were isolated from necrotic tissue at lower frequency. We did not attempt to identify fungi that were isolated at very low frequency (Table 2: Fungal isolation +).

A comparison of the rDNA-ITS region of eight *Fusarium* isolates from trees P2 and P3 revealed two patterns for the nucleotide sequences with only one nucleotide substitution (Table 2). One of the eight samples derived from whitish mycelia that grew from the surface of the discolored xylem severely damaged tree P2 (P2-fk, Table 2; Fig. 4a). Nucleotide sequence analysis of the ITS, the *RPB2*, and the *EF-1a* of representative isolate P2-f11 (strain A) indicated that strain A belongs to *F. solani* species complex (FSSC) clade 3 defined by O’Donnell et al. (2008). Figure 6 shows the phylogenetic tree inferred from the nucleotide sequences of *EF-1a* using 29 *Fusarium* strains and strain A. Strain A was adjacent to Clade B of the Ambrosia *Fusarium* clade (AFC), known as a group of symbionts of ambrosia beetles (Kasson et al. 2013; O’Donnell et al. 2015), and with *F. pseudensiforme* strains NRRL 46517 and FRC S1834, which are members of Clade A of the AFC (Kasson et al. 2013). Fungi belonging to the FSSC were not detected in the non-discolored xylem (Table 2).

Fig. 4 *Fusarium* sp. isolated from declining *E. variegata*. **a-1** Whitish fungal hyphae on cross-cut surface of *E. variegata* trunk. **a-2** Enlarged image of the boxed region in **a-1**. **b** Strain A (P2-f11). **c** Strain B (P2-g21). **d** Conidia of P2-f11. **e** Conidia of P2-g21



Quadrastichus erythrinae and fungi in leaves and petioles

Larvae of *Q. erythrinae* were found in galls on the leaves or petioles of specimens R4–R6 (with abundant leaves, Level 2). *Fusarium* spp. were isolated from some petioles of R3, R5, and R6, including normal leaves of R3 without galls.

Fungal distribution in controls without Q. erythrinae infestation

No fungi were isolated from the stems, and no *Fusarium* spp. were isolated from the petioles of control seedlings, C1–C3 (Table 2).

Xylem dysfunction of host trees tested by dye injection

In the heavily damaged tree P2, absorption of the dye solution through vessels was only 6.3 mL/h. A very narrow area was stained with the dye (Fig. 2a, Sa), since water conduction had almost ceased in the trunk by the time of the experiment. In contrast, tree P3, with abundant leaves, absorbed the dye solution at 15 mL/h, and the area with stained xylem was broader than that of P2 (Fig. 2b, Sa). These results demonstrated that water conduction in the trunk of P3 was better than in P2. However, most leaves had drooped on tree P3 by around 2 p.m., when





Fig. 5 Petiole structure of *E. variegata* with or without galls of *Quadrastichus erythrinae*. **a** Shoot with galls of *Q. erythrinae*. **b** Cross section of normal petiole tissue showing clear boundaries of cortex (cor), primary phloem (pp), primary xylem (px), and pith (p). **c** Abnormal tissue in petiole with a gall showing hyperplasia and hypertrophy of cells. Sections **b** and **c** were stained with toluidine blue O

transpiration was very active, meaning that the water supply to shoots and leaves was insufficient even in tree P3, which had abundant leaves.

Fig. 6 Maximum likelihood (ML) phylogenetic tree based on *EF-1a* nucleotide sequences. Numbers at nodes represent bootstrap support >50% from 1000 replicates. Three clades in the FSSC and the Ambrosia *Fusarium* Clade were indicated by bold internodes. Accession numbers and NRRL (the ARS Culture Collection) or FRC (Fusarium Research Center) numbers are shown



Effect of *Fusarium* spp. and *F. solani* after inoculation of seedlings

In the preliminary inoculation of healthy seedlings, isolates P1-1600A7 (strain A) and R1-BC6 (strain B), which were detected in trees P1 and R1 (Table 2), induced brownish discoloration in the xylem and necrosis in the phloem. We then inoculated the roots with strains A (P2-f11) and B (P2-f22) each one singly or with a mixture (Table 3, footnote). The mixture of strains A and B killed a seedling in 26 days, and strain A killed another seedling in 35 days (Fig. 7). The lower stems became soft (Fig. 7b, c), and at the microscopic level, cell necrosis was similar to that seen from decline of tree in the field. Whitish hyphae emerged from the necrotic area of the lower stems (Fig. 7c) and were confirmed as inoculated strain A. Seedlings inoculated with strain B did not show symptoms. On the 46th day after inoculation, November 17, one of the control seedlings died without phloem necrosis following defoliation, and strains A and B were not isolated from the dead tissue.

In the third experiment, among nine inoculated seedlings, three seedlings died during the third to sixth weeks after inoculation, and seven seedlings developed slight symptoms, such as leaf yellowing and partial defoliation, then recovered. All seedlings inoculated with strain A, dead or surviving, had xylem discoloration and necrosis of the cortex and phloem in the lower stems and an odor. Strain A




Fig. 7 *E. variegata* seedlings either non-inoculated or inoculated with a mixture of conidia of strains A and B. **a** Healthy seedling. **b** Dead seedling, 26 days after inoculation. **c** Whitish hyphae (arrow) from softened and necrotic tissue from the dead seedling in image **b**

was reisolated from all inoculated specimens. In contrast, two control seedlings inoculated with sterilized toothpicks showed no symptoms, and strain A was not isolated.

Discussion

Brown to gray discoloration of the xylem in the lower trunk or thick branch was characteristic of the severely declining, defoliated *E. variegata* trees. This type of xylem discoloration is known as wound heartwood or pathological heartwood, caused by the defense reaction against infection; in turn, water conduction ceases in this area, as with normal heartwood (Hillis 1987). Brownish discoloration of the host tissue is caused by the secretion of secondary metabolites.

In the present study, loss of water conduction in declining trees was detected via the dye injection method, used in field and inoculation experiments to visualize water flow within specimens (Kuroda 2001; Morita et al. 2016). We found that water conduction had almost ceased in the heavily damaged *E. variegata* tree, which had wide discoloration in the basal part of the trunk, in Peace Memorial Park. In the slightly damaged tree, the water supply was insufficient, as evidenced by temporary leaf drooping during the daytime and the small amount of dye absorption. These results suggest that blockage of water conduction had progressed before the defoliation.

The causal relationship between xylem discoloration and water conduction blockage in trees has been well explained in reports on wilt diseases of trees. For instance, Japanese oak wilt caused by *Raffaelea quercivora* and fig wilt (fig canker) caused by *Ceratocystis ficicola* are wilt diseases that induce severe discoloration in the xylem and the dysfunction of water conduits at the same time (Kajiji et al. 2013; Kuroda

2001; Morita et al. 2016; Sumida et al. 2016). Infected trees die from the cessation of water conduction when the host defense is ineffective and the discolored and dysfunctional area has almost covered the entire cross section of the trunks (Kuroda 2001). This phenomenon is universal in wilt diseases of broadleaved trees.

We always detected fungi that belong to the *Fusarium solani* species complex (FSSC) from heavily deteriorated or almost dead trees. Some strains, including strains A and B, might be causing the decline and death of *E. variegata* because these strains were predominantly isolated from the discolored xylem of declining trees, and fungal hyphae were observed in the same area using microscopy. The tylosis formation also indicates vessel dysfunction in the discolored areas. In addition, secretion duct occlusions in the specimens are a sign of a defense reaction, in contrast to other reports on *Erythrina* trees in which occlusions were not found (Teixeira et al. 2000; Silva et al. 2013). In the slightly damaged trees with abundant leaves, the xylem discoloration was partial and pale in color. The area with this defense reaction and discoloration seem to have expanded by harvest. The lack of isolation of fungi belonging to the FSSC from non-discolored xylem supports the idea that the defense reaction of the host progressed in association with growth of strains of *Fusarium* sp.

Phloem necrosis with an odor always occurred adjacent to the discolored xylem, and the same strains of *Fusarium* spp. were detected in both areas. Therefore, we focused on the identification of isolated strains and inoculation experiments. Other fungi, the *Papularia* sp., *Pestalotia* sp., and *Alternaria* sp., isolated with very low frequency, were judged to be a secondary infection to the necrotic host tissue and were not considered further as candidates.

In the phylogenetic analysis, strain A was a member of the Ambrosia *Fusarium* clade (AFC), especially closely related to *F. pseudensiforme*, which is a member of Clade A of the AFC (Fig. 6) and known as the causal agent of *Fusarium* dieback of avocado (Eskalen et al. 2012; Kasson et al. 2013). The AFC belongs to FSSC clade 3, which is composed of pathologically important species of both humans and plants (Aoki et al. 2014; O'Donnell et al. 2008; Zhang et al. 2006). Strain A was always isolated from the discolored xylem and necrotic phloem of *E. variegata* before tree death. Ambrosia beetles, such as the tea shot-hole borer (*Euwallacea fornicatus*), is known as a vector of *Fusarium* spp. in the AFC (Eskalen et al. 2012; Kasson et al. 2013). Detailed field research will be necessary to understand the relationships between strain A detected in *E. variegata* and the fungal symbionts of ambrosia beetles. *F. virguliforme*, a member of the FSSC clade 2, is known to cause sudden death syndrome in the soybean *Glycine max* (Roy et al. 1997), which belongs to the same family, Fabaceae, as *Erythrina*. Several reports indicate similarities between the symptoms of

soybean SDS and those of *Erythrina* decline, such as the discoloration of stem tissue and tissue necrosis (Aoki et al. 2003, 2014; Roy et al. 1997). Note that the causal agent of SDS was confirmed only after a long history of investigation because experimental demonstration was difficult (Radwan et al. 2011).

Inoculation experiments in the present study were preliminary because seedling preparation was difficult and the experiments were done in a cooler climate than in the area where natural infection occurs. Nevertheless, seedlings inoculated with strain A died after the development of microscopic symptoms similar to those observed on naturally infected trees, and the inoculated strain was reisolated from all samples from symptomatic or dead seedlings. Therefore, we judged that at least one strain of the FSSC is pathogenic on *E. variegata* seedlings. Mortality rates typically are not as high in inoculation experiments for wilt diseases of trees (Morita et al. 2016), likely because xylem dysfunction progresses gradually in the host stems. So trees do not die until several years after infection. Repeated inoculation experiments should provide more information on disease development.

This incidence is a new discovery regarding the contribution of a fungus to the decline and death of *E. variegata*. Although the phylogenetic investigation of strains A and B within the FSSC is insufficient, reporting on this incidence is important for the development of new strategies to rescue *Erythrina* spp. If *Q. erythrinae* is not responsible for the decline of *Erythrina* trees, releasing *Eurytoma erythrinae*, a wasp indigenous to Africa, to biologically control *Q. erythrinae* in Hawaii (Bell et al. 2013; Yalem et al. 2009) will be misdirected. This strategy will also need to be reconsidered in Japan's Okinawa Prefecture.

Strain A was mostly found in the lower part of severely damaged *E. variegata* trunks and in thick branches. Because *Fusarium* species mainly inhabit soil, spread of infection to adjacent trees may occur via root contact with infected roots or wounds in trunk bases, often caused by a lawn mower. Information from the prefectural government that the decline is progressing adjacent to dead trees also supports our prediction of fungal infection.

Based on the present observations, the contribution of *Q. erythrinae* to the xylem discoloration, phloem necrosis with odor, and death of *Erythrina* is doubtful. A defense reaction was not found around the galls of *Q. erythrinae*. Phloem necrosis was associated with xylem discoloration and was not observed around the galls. Detailed observation will be necessary to confirm the infection process and to determine whether *Q. erythrinae* is contributing to the decline of *Erythrina*. If it does not directly cause the tree's death, we need to assess whether the wasp is causing physiological stress to the hosts and promoting symptom development.

The slow defoliation of *E. variegata* trees over several years suggests that, although death appears to occur suddenly, xylem dysfunction may progress over many years. Infection by strain A indicated its pathogenicity to *E. variegata* seedlings; however, it may not be virulent enough to kill large trees in a short period. Perhaps a stressed condition is necessary to promote symptoms and cause death. Fungal strains should be collected from other sites in Japan and other areas where *Erythrina* decline is occurring to further examine the pathogenicity of this fungus. The distance of sampling sites, approximately 24 km between the University of the Ryukyus and Okinawa Prefectural Peace Memorial Park, suggests that the contribution of strain A to the *E. variegata* decline may be common over a wide area in Okinawa.

To conclude, *Fusarium* sp. may be a causal agent of xylem dysfunction and phloem necrosis in *E. variegata*. *Q. erythrinae* does not seem to contribute to the decline and death, since xylem discoloration and the distribution of pathogenic strain A were not found around the galls of *Q. erythrinae*. In addition, we obtained information that *Fusarium* sp. had been detected in an *E. crista-galli* specimen that died suddenly three decades ago in Kagoshima (J. Miyajima, Kumamoto Prefectural Government, personal communication), where *Q. erythrinae* was not present. These observations suggest that the sudden death of *Erythrina* occurs without *Q. erythrinae* infestation. Still, *Q. erythrinae* could promote the decline of its hosts through physiological stress. Stressed *E. variegata* may become susceptible to strain A. If the decline and death of *E. variegata* increased after the *Q. erythrinae* spread throughout the southern islands of Japan, the role of this wasp must be investigated from a pathological perspective. Although we need detailed investigations to elucidate the entire process of *Erythrinae* decline and death, information on the pathogenicity of strain A to *E. variegata* will be helpful in developing methods to protect hosts. Injecting insecticide into trunks does not stop *Erythrinae* sudden death. So an alternative treatment, fungicide injection, may prevent xylem colonization by strain A and maintain the ascent of sap. We have begun test injections of a fungicide that is popular for protecting oak trees from Japanese oak wilt.

Acknowledgements The authors are grateful to Dr. Takayuki Aoki (National Institute of Agrobiological Sciences, Genetic Resources Center) for his advice on the phylogenetic analysis of *Fusarium* spp. We also thank the Okinawa Prefectural Forest Resource Research Center and Ryukyu Sankei Co., Ltd., for supplying *E. variegata* seedlings for inoculation experiments. This work was supported by JSPS KAKENHI Grant Number JP15K14747 and by Okinawa Prefecture.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give

appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Aoki T, O'Donnell K, Homma Y, Lattanzi AR (2003) Sudden-death syndrome of soybean is caused by two morphologically and phylogenetically distinct species within the *Fusarium solani* species complex—*F. virguliforme* in North America and *F. tucumaniae* in South America. *Mycologia* 95:660–684
- Aoki T, O'Donnell K, Geiser DM (2014) Systematics of key phytopathogenic *Fusarium* species: current status and future challenges. *J Gen Plant Pathol* 80:189–201
- Barnett HL, Hunter BB (1998) Illustrated genera of imperfect fungi, 4th edn. APS Press, St. Paul, p. 218
- Bell RC, Belmaker A, Couch CS, Marchetto KM, Simonis JL, Thomas RQ, Spark JP, Brown JM, Francisco KS, Manuel ME (2013) Effectiveness of *Erythrina* gall wasp biocontrol and implications for the recovery of threatened Wiliwili trees (Fabaceae: *Erythrina sandwicensis*). *J Torrey Bot Soc* 140:215–224
- Costa SS, Matos KS, Tessmann DJ, Seixas CDS, Pfenning LH (2016) *Fusarium paranaense* sp. nov., a member of the *Fusarium solani* species complex causes root rot on soybean in Brazil. *Fungal Biol* 120:51–60
- Culik MP, Martins DDS, Ventura JA, Costa VA (2014) The invasive gall wasp *Quadrastichus erythrinae* (Hymenoptera: Eulophidae) in South America: is classical biological control needed? *Biocontrol Sci Techn* 24(8):971–975
- Eskalen A, Gonzalez A, Wang DH, Twizeyimana M, Mayorquin JS, Lynch SC (2012) First report of a *Fusarium* sp. and its vector tea shot hole borer (*Euwallacea fornicatus*) causing *Fusarium* dieback on avocado in California. *Plant Dis* 96:1070
- Faizal MH, Prathapan KD, Anith KN, Mary CA, Lekha M, Rini CR (2006) *Erythrina* gall wasp *Quadrastichus erythrinae*, yet another invasive pest new to India. *Curr Sci* 90:1061–1062
- Feder N, O'Brien TP (1968) Plant microtechnique: some principles and new methods. *Am J Bot* 55:123–139
- Gramling C (2005) Hawaii's coral trees feel the sting of foreign wasps. *Science* 310:1759–1760
- Heu RA, Tsuda DM, Nagamine WT, Yalamar JA, Suh TH (2008) *Erythrina* gall wasp *Quadrastichus erythrinae* Kim (Hymenoptera: Eulophidae). New Pest Advisory No. 05-03, Hawaii Department of Agriculture, Honolulu
- Hillis WE (1987) Heartwood and tree exudates. Springer-Verlag, Berlin, pp 157–162
- Jukes TH, Cantor CR (1969) Evolution of protein molecules. In: Munro HN (ed) Mammalian protein metabolism. Academic Press, New York, pp 21–132
- Kajii C, Morita T, Jikumaru S, Kajimura H, Yamaoka Y, Kuroda K (2013) Xylem dysfunction in *Ficus carica* infected with wilt fungus *Ceratocystis ficicola* and the role of the vector beetle *Euwallacea interjectus*. *IAWA J* 34:301–312
- Kasson MT, O'Donnell K, Rooney AP, Sink S, Ploetz RC, Ploetz JN, Konkol JL, Carrillo D, Freeman S, Mendel Z, Smith JA, Black AW, Hulcr J, Bateman C, Stefkova K, Campbell PR, Geering AD, Dann EK, Eskalen A, Mohotti K, Short DP, Aoki T, Fenstermacher KA, Davis DD, Geiser DM (2013) An inordinate fondness for *Fusarium*: phylogenetic diversity of fusaria cultivated by ambrosia beetles in the genus *Euwallacea* on avocado and other plant hosts. *Fungal Genet Biol* 56:147–157
- Kihara T, Murakami T, Chuma I, Kameyama N, Kuroda K (2016) A pathological and anatomical study on the factor of phloem necrosis and wilt of *Erythrina variegata* (in Japanese). *Tree For-est Health* 20:95–96
- Kim IK, Delvare G, La Salle J (2004) A new species of *Quadrastichus* (Hymenoptera: Eulophidae): a gall-inducing pest on *Erythrina* (Fabaceae). *J Hymenoptera Res* 13(2):243–249
- Kiyuna C (2008) Effect of a pesticide injection to *Erythrina variegata* to kill *Quadrastichus erythrinae* (in Japanese). *Res Rep Okinawa Prefect For Resour Res Cent* 50:10–1
- Kumar S, Stecher G, Tamura K (2016) MEGA7: Molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Mol Biol Evol* 33:1870–1874
- Kuroda K (2001) Responses of *Quercus* sapwood to infection with the pathogenic fungus of a new wilt disease vectored by the ambrosia beetle *Platypus quercivorus*. *J Wood Sci* 47:425–429
- Kuroda K (2005) Xylem dysfunction in Yezo spruce (*Picea jezoensis*) after inoculation with the blue-stain fungus *Ceratocystis polonica*. *For Pathol* 35:346–358
- Kuroda K, Yamada T (1996) Discoloration of sapwood and blockage of xylem sap ascent in the trunks of wilting *Quercus* spp. following attack by *Platypus quercivorus* (in Japanese with English summary). *J Jpn For Soc* 78:84–88
- Morita T, Jikumaru S, Kuroda K (2016) Disease development in *Ficus carica* plants after inoculation with *Ceratocystis ficicola*. (1) relationships between xylem dysfunction and wilt symptom (in Japanese with English summary). *Jpn J Phytopathol* 82:301–309
- O'Donnell K, Sarver BA, Brandt M, Chang DC, Noble-Wang J, Park BJ, Sutton DA, Benjamin L, Lindsley M, Padhye A, Geiser DM, Ward TJ (2007) Phylogenetic diversity and microsphere array-based genotyping of human pathogenic *Fusaria*, including isolates from the multistate contact lens-associated U.S. keratitis outbreaks of 2005 and 2006. *J Clin Microbiol* 45:2235–2248
- O'Donnell K (2000) Molecular phylogeny of the *Nectria haematococca*–*Fusarium solani* species complex. *Mycologia* 92:919–938
- O'Donnell K, Sutton DA, Fothergill A, McCarthy D, Rinaldi MG, Brandt ME, Zhang N, Geiser DM (2008) Molecular phylogenetic diversity, multilocus haplotype nomenclature, and in vitro antifungal resistance within the *Fusarium solani* species complex. *J Clin Microbiol* 46:2477–2490
- O'Donnell K, Sink S, Libeskind-Hadas R, Hulcr J, Kasson MT, Ploetz RC, Konkol JL, Ploetz JN, Carrillo D, Campbell A, Duncan RE, Liyanage PNH, Eskalen A, Na F, Geiser DM, Bateman C, Freeman S, Mendel Z, Sharon M, Aoki T, Cossé AA, Rooney AP (2015) Discordant phylogenies suggest repeated host shifts in the *Fusarium*–*Euwallacea* ambrosia beetle mutualism. *Fungal Genet Biol* 82:277–290
- Radwan O, Liu Y, Clough SJ (2011) Transcriptional analysis of soybean root response to *Fusarium virguliforme*, the causal agent of sudden death syndrome. *Mol Plant-Microbe Interact* 24:958–972
- Reimar NJ (2007) Field release of *Eurytoma* sp. (Hymenoptera: Eurytomidae), for biological control of the *Erythrina* gall wasp, *Quadrastichus erythrinae* Kim (Hymenoptera: Eulophidae), in Hawaii. Draft Environmental Assessment, p 8
- Roy KW, Rupe JC, Hershman DE, Abney TS (1997) Sudden death syndrome of soybean. *Plant Dis* 81:1100–1111
- Rubinoff D, Holland BS, Shibata A, Messing RH, Wright MG (2010) Rapid invasion despite lack of genetic variation in the *Erythrina* gall wasp (*Quadrastichus erythrinae* Kim). *Pacific Sci* 64:23–31
- Ruzin SE (1999) Plant microtechnique and microscopy. Oxford University, New York, pp 87–95
- Silva MB da, Santana ASCO, Pimentel RMM, Silva FCL, Randau KP, Soares LAL (2013) Anatomy of leaf and stem of *Erythrina velutina*. *Rev Bras Farmacogn* 23:200–206
- Sumida S, Kajii C, Morita T, Kuroda K (2016) Disease development in *Ficus carica* seedlings after inoculation with *Ceratocystis ficicola*. (2) microscopic analysis of the host-pathogen interaction and internal-

- symptoms (Japanese with English summary). *Jpn J Phytopathol* 82:310–316
- Teixeira SDP, Castro MDM, Tozzi ADA (2000) Secretory cavities and pellucid dots in leaflets of *Lonchocarpus* (Leguminosae, Papilionoideae, Millettieae). *Plant System Evol* 221:61–68
- Uechi N, Uesato T, Yukawa J (2007) Detection of an invasive gall-inducing pest, *Quadrastichus erythrinae* (Hymenoptera: Eulophidae), causing damage to *Erythrina variegata* L. (Fabaceae) in Okinawa Prefecture, Japan. *Entomol Sci* 10:209–212
- White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ (eds.) *PCR protocols: a guide to methods and applications*. Academic Press, San Diego, pp 315–322
- Yalemar J, Nagamine W, Ramadan M, Bautista R (2009) Managing the *Erythrina* gall wasp problem in Hawaii by classical biological control. Presentation of State of Hawaii Department of Agriculture, Honolulu, p. 27
- Yang MM, Tung GS, LaSalle J, Wu ML (2004) Outbreak of erythrina gall wasp (Hymenoptera: Eulophidae) on *Erythrina* spp. (Fabaceae). Taiwan. *Plant Protect Bull* 46:391–396
- Zhang N, O'Donnell K, Sutton DA, Nalim FA, Summerbell RC, Padhye AA, Geiser DM (2006) Members of the *Fusarium solani* species complex that cause infections in both humans and plants are common in the environment. *J Clin Microbiol* 44:2186–2190