

Hitchin's system and the Geometric Langlands conjecture

Ron Donagi

Penn

Kobe, December 7, 2016

Introduction

This talk has two goals.

Introduction

This talk has two goals.

The first goal of this talk is to help celebrate Saito-San's birthday.

Introduction

This talk has two goals.

The first goal of this talk is to help celebrate Saito-San's birthday.



Introduction

This talk has two goals.

The first goal of this talk is to help celebrate Saito-San's birthday

Introduction

This talk has two goals.

The first goal of this talk is to help celebrate Saito-San's birthday and his many contributions to integrable systems and moduli spaces of connections.

Introduction

This talk has two goals.

The first goal of this talk is to help celebrate Saito-San's birthday and his many contributions to integrable systems and moduli spaces of connections.

The second goal of this talk is to explain some aspects of the Geometric Langlands Conjecture, as it relates to:

Introduction

This talk has two goals.

The first goal of this talk is to help celebrate Saito-San's birthday and his many contributions to integrable systems and moduli spaces of connections.

The second goal of this talk is to explain some aspects of the Geometric Langlands Conjecture, as it relates to:

- (homological mirror symmetry)

Introduction

This talk has two goals.

The first goal of this talk is to help celebrate Saito-San's birthday and his many contributions to integrable systems and moduli spaces of connections.

The second goal of this talk is to explain some aspects of the Geometric Langlands Conjecture, as it relates to:

- (homological mirror symmetry)
- Hitchin's system, and

Introduction

This talk has two goals.

The first goal of this talk is to help celebrate Saito-San's birthday and his many contributions to integrable systems and moduli spaces of connections.

The second goal of this talk is to explain some aspects of the Geometric Langlands Conjecture, as it relates to:

- (homological mirror symmetry)
- Hitchin's system, and
- non abelian Hodge theory.

Introduction

This talk has two goals.

The first goal of this talk is to help celebrate Saito-San's birthday and his many contributions to integrable systems and moduli spaces of connections.

The second goal of this talk is to explain some aspects of the Geometric Langlands Conjecture, as it relates to:

- (homological mirror symmetry)
- Hitchin's system, and
- non abelian Hodge theory.

Joint work with Tony Pantev, and partially with Carlos Simpson.

Geometric Langlands

The conjecture:

A G -local system \mathbb{V} on a Riemann surface $C \Leftrightarrow$
an automorphic D -module $D_{\mathbb{V}}$ on the moduli space $\mathbf{Bun}_{C, {}^L G}$ of
 ${}^L G$ -bundles on C .

Geometric Langlands

The conjecture:

A G -local system \mathbb{V} on a Riemann surface $C \Leftrightarrow$
an automorphic D -module $D_{\mathbb{V}}$ on the moduli space $\mathbf{Bun}_{C, {}^L G}$ of
 ${}^L G$ -bundles on C .
(${}^L G$ is the Langlands dual group of G .)

Geometric Langlands

The conjecture:

A G -local system \mathbb{V} on a Riemann surface $C \Leftrightarrow$
an automorphic D -module $D_{\mathbb{V}}$ on the moduli space $\mathbf{Bun}_{C, {}^L G}$ of
 ${}^L G$ -bundles on C .

(${}^L G$ is the Langlands dual group of G . It is characterized by
 $\text{cochar}_{[{}^L G]} = \text{char}_{[G]}$.)

Geometric Langlands

The conjecture:

A G -local system \mathbb{V} on a Riemann surface $C \Leftrightarrow$
an automorphic D -module $\mathcal{O}_{\mathbb{V}}$ on the moduli space $\mathbf{Bun}_{C, {}^L G}$ of
 ${}^L G$ -bundles on C .

Geometric Langlands

The conjecture:

A G -local system \mathbb{V} on a Riemann surface $C \Leftrightarrow$
 an automorphic D -module $c_{\mathbb{V}}$ on the moduli space $\mathbf{Bun}_{C, L^1 G}$ of
 $L^1 G$ -bundles on C .

A fancier version of the Geometric Langlands conjecture predicts the
 existence of a canonical equivalence of categories

$$\mathfrak{c} : D_{\text{coh}}(\mathbf{Loc}, \mathcal{O}) \xrightarrow{\cong} D_{\text{coh}}({}^L \mathbf{Bun}, \mathcal{D}), \quad (\text{GLC})$$

which is uniquely characterized by the property that \mathfrak{c} sends the
 structure sheaves of points \mathbb{V} in \mathbf{Loc} to Hecke eigen \mathcal{D} -modules $\mathfrak{c}(\mathcal{O}_{\mathbb{V}})$
 (corresponding to the above $c_{\mathbb{V}}$) on ${}^L \mathbf{Bun}$:

$${}^L H^{\mu}(\mathfrak{c}(\mathcal{O}_{\mathbb{V}})) = \mathfrak{c}(\mathcal{O}_{\mathbb{V}}) \boxtimes \rho^{\mu}(\mathbb{V}).$$

Geometric Langlands

The conjecture:

A G -local system \mathbb{V} on a Riemann surface $C \Leftrightarrow$
 an automorphic \mathcal{D} -module $c_{\mathbb{V}}$ on the moduli space $\mathbf{Bun}_{C, {}^L G}$ of
 ${}^L G$ -bundles on C .

A fancier version of the Geometric Langlands conjecture predicts the
 existence of a canonical equivalence of categories

$$\mathfrak{c} : D_{\text{coh}}(\mathbf{Loc}, \mathcal{O}) \xrightarrow{\cong} D_{\text{coh}}({}^L \mathbf{Bun}, \mathcal{D}), \quad (\text{GLC})$$

which is uniquely characterized by the property that \mathfrak{c} sends the
 structure sheaves of points \mathbb{V} in \mathbf{Loc} to Hecke eigen \mathcal{D} -modules $\mathfrak{c}(\mathcal{O}_{\mathbb{V}})$
 (corresponding to the above $c_{\mathbb{V}}$) on ${}^L \mathbf{Bun}$:

$${}^L H^{\mu}(\mathfrak{c}(\mathcal{O}_{\mathbb{V}})) = \mathfrak{c}(\mathcal{O}_{\mathbb{V}}) \boxtimes \rho^{\mu}(\mathbb{V}).$$

Here μ is an appropriate character, and ${}^L H^{\mu}$ is the Hecke
 correspondence bounded by μ .

Geometric Langlands

Motivation:

The Langlands program is the *non-abelian* extension of class field theory (CFT). The abelian case is well understood. Its geometric version, or geometric CFT, is essentially the theory of a curve C and its Jacobian $J = J(C)$. This abelian case of the Geometric Langlands Conjecture (GLC) amounts to the well known result that any rank 1 local system (or: line bundle with flat connection) on the curve C extends uniquely to J , and this extension is natural with respect to the Abel-Jacobi map. The structure group of a rank 1 local system is of course just the abelian group $C^* = GL(1, C)$. The GLC is the attempt to extend this classical result from C^* to all complex reductive groups G . This goes as follows.

Geometric Langlands

The **Jacobian** is replaced by the moduli stack **Bun** of principal bundles V on C whose structure group is the Langlands dual group ${}^L G$ of the original G .

Geometric Langlands

The **Jacobian** is replaced by the moduli stack **Bun** of principal bundles V on C whose structure group is the Langlands dual group ${}^L G$ of the original G . The analogues of the **Abel-Jacobi maps** are the *Hecke correspondences* $H \subset \mathbf{Bun} \times \mathbf{Bun} \times C$.

Geometric Langlands

The **Jacobian** is replaced by the moduli stack **Bun** of principal bundles V on C whose structure group is the Langlands dual group ${}^L G$ of the original G . The analogues of the **Abel-Jacobi maps** are the *Hecke correspondences* $H \subset \mathbf{Bun} \times \mathbf{Bun} \times C$. These parametrize quadruples (V, V', x, β) where x is a point of C , while V, V' are $({}^L G)$ -bundles on C , with an isomorphism $\beta : V|_{C-x} \rightarrow V'|_{C-x}$ away from the point x having prescribed order of blowing up at x .

Geometric Langlands

The **Jacobian** is replaced by the moduli stack **Bun** of principal bundles V on C whose structure group is the Langlands dual group ${}^L G$ of the original G . The analogues of the **Abel-Jacobi maps** are the *Hecke correspondences* $H \subset \mathbf{Bun} \times \mathbf{Bun} \times C$. These parametrize quadruples (V, V', x, β) where x is a point of C , while V, V' are $({}^L G)$ -bundles on C , with an isomorphism $\beta : V|_{C-x} \rightarrow V'|_{C-x}$ away from the point x having prescribed order of blowing up at x . (In case $G = {}^L G = C^*$ these become triples (L, L', x) where the line bundle L' is obtained from L by tensoring with some fixed power of the line bundle $\mathcal{O}_C(x)$. By fixing L and varying x we see that this is indeed essentially the Abel-Jacobi map.)

Geometric Langlands

The **Jacobian** is replaced by the moduli stack **Bun** of principal bundles V on C whose structure group is the Langlands dual group ${}^L G$ of the original G . The analogues of the **Abel-Jacobi maps** are the *Hecke correspondences* $H \subset \mathbf{Bun} \times \mathbf{Bun} \times C$. These parametrize quadruples (V, V', x, β) where x is a point of C , while V, V' are $({}^L G)$ -bundles on C , with an isomorphism $\beta : V|_{C-x} \rightarrow V'|_{C-x}$ away from the point x having prescribed order of blowing up at x . (In case $G = {}^L G = C^*$ these become triples (L, L', x) where the line bundle L' is obtained from L by tensoring with some fixed power of the line bundle $\mathcal{O}_C(x)$. By fixing L and varying x we see that this is indeed essentially the Abel-Jacobi map.) For bigger groups, there are many ways to specify the allowed order of growth of β , indexed by dominant characters $\mu \in \text{char}^+_{[{}^L G]} = \text{cochar}^+_{[G]}$. So there is a collection of Hecke correspondences H^μ , each inducing a Hecke operator on various categories of objects on **Bun**.

Geometric Langlands

The **Jacobian** is replaced by the moduli stack **Bun** of principal bundles V on C whose structure group is the Langlands dual group ${}^L G$ of the original G . The analogues of the **Abel-Jacobi maps** are the *Hecke correspondences* $H \subset \mathbf{Bun} \times \mathbf{Bun} \times C$. These parametrize quadruples (V, V', x, β) where x is a point of C , while V, V' are $({}^L G)$ -bundles on C , with an isomorphism $\beta : V|_{C-x} \rightarrow V'|_{C-x}$ away from the point x having prescribed order of blowing up at x . (In case $G = {}^L G = C^*$ these become triples (L, L', x) where the line bundle L' is obtained from L by tensoring with some fixed power of the line bundle $\mathcal{O}_C(x)$. By fixing L and varying x we see that this is indeed essentially the Abel-Jacobi map.) For bigger groups, there are many ways to specify the allowed order of growth of β , indexed by dominant characters $\mu \in \text{char}^+_{[{}^L G]} = \text{cochar}^+_{[G]}$. So there is a collection of Hecke correspondences H^μ , each inducing a Hecke operator on various categories of objects on **Bun**. The resulting Hecke operators form a **commutative algebra**

Geometric Langlands

The **Jacobian** is replaced by the moduli stack **Bun** of principal bundles V on C whose structure group is the Langlands dual group ${}^L G$ of the original G . The analogues of the **Abel-Jacobi maps** are the *Hecke correspondences* $H \subset \mathbf{Bun} \times \mathbf{Bun} \times C$. These parametrize quadruples (V, V', x, β) where x is a point of C , while V, V' are $({}^L G)$ -bundles on C , with an isomorphism $\beta : V|_{C-x} \rightarrow V'|_{C-x}$ away from the point x having prescribed order of blowing up at x . (In case $G = {}^L G = C^*$ these become triples (L, L', x) where the line bundle L' is obtained from L by tensoring with some fixed power of the line bundle $\mathcal{O}_C(x)$. By fixing L and varying x we see that this is indeed essentially the Abel-Jacobi map.) For bigger groups, there are many ways to specify the allowed order of growth of β , indexed by dominant characters $\mu \in \text{char}^+_{[{}^L G]} = \text{cochar}^+_{[G]}$. So there is a collection of Hecke correspondences H^μ , each inducing a Hecke operator on various categories of objects on **Bun**. The resulting Hecke operators form a **commutative algebra**, so can have simultaneous eigen-objects.

Correspondences

A correspondence between varieties A, B is a subscheme $H \subset A \times B$. It induces a transform from objects on A to objects on B , by pull-push. (May have to specify a kernel.) Each Hecke correspondence $H^\mu \subset \mathbf{Bun} \times \mathbf{Bun} \times C$ induces a transform from D-modules on \mathbf{Bun} to D-modules on $\mathbf{Bun} \times C$.

Geometric Langlands

The Geometric Langlands Conjecture says that an irreducible G -local system on C determines a perverse sheaf on \mathbf{Bun} which is a simultaneous eigensheaf for the action of the Hecke operators - this turns out to be the right generalization of naturality with respect to the Abel-Jacobi map. (A perverse sheaf is, roughly, a local system on a Zariski open subset of \mathbf{Bun} , extended in a natural way across the complement.) Fancier versions of the conjecture recast this as an equivalence of derived categories: of D -modules on \mathbf{Bun} vs. coherent sheaves on the moduli space \mathbf{Loc} of local systems. There are many related conjectures and extensions, notably to punctured curves via parabolic bundles and local systems. Some of these make an appearance below.

Geometric Langlands

The conjecture:

A G -local system \mathbb{V} on a Riemann surface $C \Leftrightarrow$
 an automorphic \mathcal{D} -module $c_{\mathbb{V}}$ on the moduli space $\mathbf{Bun}_{C, L^1 G}$ of
 $L^1 G$ -bundles on C .

A fancier version of the Geometric Langlands conjecture predicts the
 existence of a canonical equivalence of categories

$$\mathfrak{c} : D_{\text{coh}}(\mathbf{Loc}, \mathcal{O}) \xrightarrow{\cong} D_{\text{coh}}(L^1 \mathbf{Bun}, \mathcal{D}), \quad (\text{GLC})$$

which is uniquely characterized by the property that \mathfrak{c} sends the
 structure sheaves of points \mathbb{V} in \mathbf{Loc} to Hecke eigen \mathcal{D} -modules $\mathfrak{c}(\mathcal{O}_{\mathbb{V}})$
 (corresponding to the above $c_{\mathbb{V}}$) on $L^1 \mathbf{Bun}$:

$${}^L H^{\mu}(\mathfrak{c}(\mathcal{O}_{\mathbb{V}})) = \mathfrak{c}(\mathcal{O}_{\mathbb{V}}) \boxtimes \rho^{\mu}(\mathbb{V}).$$

Here μ is an appropriate character, and ${}^L H^{\mu}$ is the Hecke
 correspondence bounded by μ .

Some known results

Significant progress has been made towards understanding these conjectures, through works of Drinfeld, Laumon, Beilinson, Lafforgue, Frenkel, Gaitsgory, Vilonen, Heinloth, ... Some versions are known for $GL(n)$. The conjecture is unknown for other groups. There are more recent results of Heinloth in the parabolic case. Even for $GL(n)$, the proof is indirect: no construction of non-abelian Hecke eigensheaves is known.

HMS and Geometric Langlands

Idea: Hitchin's integrable system $\mathbf{Higgs}_{C,G}$ associated to (C, G) is a twistor rotation of $\mathbf{Loc}_{C,G}$. It is a hyper Kahler SYZ fibration. It also is (or: contains) $T^*\mathbf{Bun}_{C,G}$.

HMS and Geometric Langlands

Idea: Hitchin's integrable system $\mathbf{Higgs}_{C,G}$ associated to (C, G) is a twistor rotation of $\mathbf{Loc}_{C,G}$. It is a hyper Kahler SYZ fibration. It also is (or: contains) $T^*\mathbf{Bun}_{C,G}$.

HMS applied to Hitchin's system \implies

D_L should be the relative Floer homology between two Lagrangians in $\mathbf{Higgs}_{C,G^\vee}$, one fixed, the other moving:

- View L as a point of $\mathbf{Higgs}_{C,G}$. Its Fourier-Mukai dual is a Lagrangian (with line bundle) in $\mathbf{Higgs}_{C,G^\vee}$.
- A general cotangent fiber of $\mathbf{Higgs}_{C,G^\vee}$.

Abelianization via Hitchin's system

The work described here is based on an abelianization of the GLC in terms of Higgs bundles.

Abelianization via Hitchin's system

The work described here is based on an abelianization of the GLC in terms of Higgs bundles. Higgs bundles and operations on them have already been discussed in detail in Simpson's talk. But let me review briefly.

Abelianization via Hitchin's system

The work described here is based on an abelianization of the GLC in terms of Higgs bundles.

Abelianization via Hitchin's system

The work described here is based on an abelianization of the GLC in terms of Higgs bundles. A Higgs bundle is a pair (V, ϕ) consisting of a vector bundle V with a K_C -valued endomorphism $\phi : V \rightarrow V \otimes \Omega_C^1$, where $\Omega_C^1 = K_C$ is the canonical bundle of C .

Abelianization via Hitchin's system

The work described here is based on an abelianization of the GLC in terms of Higgs bundles. A Higgs bundle is a pair (V, ϕ) consisting of a vector bundle V with a K_C -valued endomorphism $\phi : V \rightarrow V \otimes \Omega_C^1$, where $\Omega_C^1 = K_C$ is the canonical bundle of C . More generally, a G -Higgs bundle is a pair (V, ϕ) consisting of a principal G -bundle V with a section ϕ of $ad(V) \otimes \Omega_C^1$, where $ad(V)$ is the adjoint vector bundle of V .

Abelianization via Hitchin's system

The work described here is based on an abelianization of the GLC in terms of Higgs bundles. A Higgs bundle is a pair (V, ϕ) consisting of a vector bundle V with a K_C -valued endomorphism $\phi : V \rightarrow V \otimes \Omega_C^1$, where $\Omega_C^1 = K_C$ is the canonical bundle of C . More generally, a G -Higgs bundle is a pair (V, ϕ) consisting of a principal G -bundle V with a section ϕ of $ad(V) \otimes \Omega_C^1$, where $ad(V)$ is the adjoint vector bundle of V . Hitchin studied the moduli space **Higgs** of such Higgs bundles (subject to an appropriate stability condition) and showed that it is an algebraically integrable system: it is algebraically symplectic, and it admits a natural map $h : \mathbf{Higgs} \rightarrow B$ to a vector space B such that the fibers are Lagrangian subvarieties. In fact the fiber over a general point $b \in B$ (in the complement of the *discriminant hypersurface*) is an abelian variety, obtained as Jacobian or Prym of an appropriate spectral cover C_b .

Abelianization via Hitchin's system

The work described here is based on an abelianization of the GLC in terms of Higgs bundles. A Higgs bundle is a pair (V, ϕ) consisting of a vector bundle V with a K_C -valued endomorphism $\phi : V \rightarrow V \otimes \Omega_C^1$, where $\Omega_C^1 = K_C$ is the canonical bundle of C . More generally, a G -Higgs bundle is a pair (V, ϕ) consisting of a principal G -bundle V with a section ϕ of $ad(V) \otimes \Omega_C^1$, where $ad(V)$ is the adjoint vector bundle of V . Hitchin studied the moduli space **Higgs** of such Higgs bundles (subject to an appropriate stability condition) and showed that it is an algebraically integrable system: it is algebraically symplectic, and it admits a natural map $h : \mathbf{Higgs} \rightarrow B$ to a vector space B such that the fibers are Lagrangian subvarieties. In fact the fiber over a general point $b \in B$ (in the complement of the *discriminant hypersurface*) is an abelian variety, obtained as Jacobian or Prym of an appropriate spectral cover C_b . The description in terms of spectral covers is somewhat ad hoc, in that it depends on the choice of a representation of the group G . A uniform description is given in terms of generalized Pryms of *cameral covers*.

Abelianization via Hitchin's system

The main result of [DP1] is formulated as a duality of the Hitchin system: There is a canonical isomorphism between the bases $B, {}^L B$ of the Hitchin system for the group G and its Langlands dual ${}^L G$, taking the discriminant in one to the discriminant in the other. Away from the discriminants, the corresponding fibers are abelian varieties, and we exhibit a canonical duality between them.

Abelianization via Hitchin's system

The main result of [DP1] is formulated as a duality of the Hitchin system: There is a canonical isomorphism between the bases $B, {}^L B$ of the Hitchin system for the group G and its Langlands dual ${}^L G$, taking the discriminant in one to the discriminant in the other. Away from the discriminants, the corresponding fibers are abelian varieties, and we exhibit a canonical duality between them. The case of the groups $GL(n)$, $SL(n)$ and $PGL(n)$ had appeared earlier in work of Hausel and Thaddeus in the context of hyperkahler mirror symmetry.

Abelianization via Hitchin's system

The main result of [DP1] is formulated as a duality of the Hitchin system: There is a canonical isomorphism between the bases $B, {}^L B$ of the Hitchin system for the group G and its Langlands dual ${}^L G$, taking the discriminant in one to the discriminant in the other. Away from the discriminants, the corresponding fibers are abelian varieties, and we exhibit a canonical duality between them. The case of the groups $GL(n)$, $SL(n)$ and $PGL(n)$ had appeared earlier in work of Hausel and Thaddeus in the context of hyperkahler mirror symmetry. There are abelianized versions of the Hecke correspondences.

Abelianization via Hitchin's system

The main result of [DP1] is formulated as a duality of the Hitchin system: There is a canonical isomorphism between the bases $B, {}^L B$ of the Hitchin system for the group G and its Langlands dual ${}^L G$, taking the discriminant in one to the discriminant in the other. Away from the discriminants, the corresponding fibers are abelian varieties, and we exhibit a canonical duality between them. The case of the groups $GL(n)$, $SL(n)$ and $PGL(n)$ had appeared earlier in work of Hausel and Thaddeus in the context of hyperkahler mirror symmetry. There are abelianized versions of the Hecke correspondences. The [DP] results allow the construction of eigenseaves for these abelianized Hecke correspondences.

Abelianized Hecke eigensheaves

On **Higgs** there are two Lagrangian fibrations:

Abelianized Hecke eigensheaves

On **Higgs** there are two Lagrangian fibrations:

The Hitchin map: $h : \mathbf{Higgs} \rightarrow B$

The projection: $\pi : \mathbf{Higgs} \supset T^* \mathbf{Bun} \rightarrow \mathbf{Bun}$

Abelianized Hecke eigensheaves

On **Higgs** there are two Lagrangian fibrations:

The Hitchin map: $h : \mathbf{Higgs} \rightarrow B$

The projection: $\pi : \mathbf{Higgs} \supset T^* \mathbf{Bun} \rightarrow \mathbf{Bun}$

(This is a rational map)

Abelianized Hecke eigensheaves

On **Higgs** there are two Lagrangian fibrations:

The Hitchin map: $h : \mathbf{Higgs} \rightarrow B$

The projection: $\pi : \mathbf{Higgs} \supset T^* \mathbf{Bun} \rightarrow \mathbf{Bun}$

(This is a rational map)

Hecke: $H \subset \mathbf{Bun} \times \mathbf{Bun} \times C$

Abelianized Hecke eigensheaves

On **Higgs** there are two Lagrangian fibrations:

The Hitchin map: $h : \mathbf{Higgs} \rightarrow B$

The projection: $\pi : \mathbf{Higgs} \supset T^* \mathbf{Bun} \rightarrow \mathbf{Bun}$

(This is a rational map)

Hecke: $H \subset \mathbf{Bun} \times \mathbf{Bun} \times C$

Abelianized Hecke: $\tilde{H} \subset \mathbf{Higgs} \times \mathbf{Higgs} \times C$

Abelianized Hecke eigensheaves

On **Higgs** there are two Lagrangian fibrations:

The Hitchin map: $h : \mathbf{Higgs} \rightarrow B$

The projection: $\pi : \mathbf{Higgs} \supset T^* \mathbf{Bun} \rightarrow \mathbf{Bun}$

(This is a rational map)

Hecke: $H \subset \mathbf{Bun} \times \mathbf{Bun} \times C$

Abelianized Hecke: $\tilde{H} \subset \mathbf{Higgs} \times \mathbf{Higgs} \times C$

Abelianized Hecke eigensheaves: π_* of degree-0 line bundles on Hitchin fibers $h^{-1}(b)$.

Abelianized Hecke eigensheaves

On **Higgs** there are two Lagrangian fibrations:

The Hitchin map: $h : \mathbf{Higgs} \rightarrow B$

The projection: $\pi : \mathbf{Higgs} \supset T^* \mathbf{Bun} \rightarrow \mathbf{Bun}$

(This is a rational map)

Hecke: $H \subset \mathbf{Bun} \times \mathbf{Bun} \times C$

Abelianized Hecke: $\tilde{H} \subset \mathbf{Higgs} \times \mathbf{Higgs} \times C$

Abelianized Hecke eigensheaves: π_* of degree-0 line bundles on Hitchin fibers $h^{-1}(b)$. We are thinking of **Bun** as the base space and of $h^{-1}(b)$ as its spectral cover.

Abelianization via Hitchin's system

It is very tempting to try to understand the relationship of this abelianized result to the full GLC. The view of the GLC pursued in [BeDr] is that it is a quantum theory. The emphasis in [BeDr] is therefore on quantizing Hitchin's system, which leads to the investigation of *opers*. One possibility, discussed in [DP1] and [Ar], is to view the full GLC as a quantum statement whose classical limit is the result in [DP1]. The idea then would be to try to prove GLC by deforming both sides of the result of [DP1] to higher and higher orders. Arinkin has carried out some deep work in this direction. But there is another path.

NAHT and GLC

I want to explore the tantalizing possibility that the abelianized version of GLC is in fact equivalent, via recent breakthroughs in Non Abelian Hodge Theory (NAHT), to the full original (non-abelian) GLC, not only to its 0-th order or classical approximation.

NAHT and GLC

I want to explore the tantalizing possibility that the abelianized version of GLC is in fact equivalent, via recent breakthroughs in Non Abelian Hodge Theory (NAHT), to the full original (non-abelian) GLC, not only to its 0-th order or classical approximation.

Instead of viewing the solution constructed in [DP1] as a classical limit of the full solution, it is interpreted as the $z = 0$ incarnation of an equivalent twistor-type object.

NAHT and GLC

I want to explore the tantalizing possibility that the abelianized version of GLC is in fact equivalent, via recent breakthroughs in Non Abelian Hodge Theory (NAHT), to the full original (non-abelian) GLC, not only to its 0-th order or classical approximation.

Instead of viewing the solution constructed in [DP1] as a classical limit of the full solution, it is interpreted as the $z = 0$ incarnation of an equivalent twistor-type object. The twistor space is fibered over the complex z -line (or its compactification).

NAHT and GLC

I want to explore the tantalizing possibility that the abelianized version of GLC is in fact equivalent, via recent breakthroughs in Non Abelian Hodge Theory (NAHT), to the full original (non-abelian) GLC, not only to its 0-th order or classical approximation.

Instead of viewing the solution constructed in [DP1] as a classical limit of the full solution, it is interpreted as the $z = 0$ incarnation of an equivalent twistor-type object. The twistor space is fibered over the complex z -line (or its compactification). (Also known as as Deligne's λ line.)

NAHT and GLC

I want to explore the tantalizing possibility that the abelianized version of GLC is in fact equivalent, via recent breakthroughs in Non Abelian Hodge Theory (NAHT), to the full original (non-abelian) GLC, not only to its 0-th order or classical approximation.

Instead of viewing the solution constructed in [DP1] as a classical limit of the full solution, it is interpreted as the $z = 0$ incarnation of an equivalent twistor-type object. The twistor space is fibered over the complex z -line (or its compactification).

NAHT and GLC

I want to explore the tantalizing possibility that the abelianized version of GLC is in fact equivalent, via recent breakthroughs in Non Abelian Hodge Theory (NAHT), to the full original (non-abelian) GLC, not only to its 0-th order or classical approximation.

Instead of viewing the solution constructed in [DP1] as a classical limit of the full solution, it is interpreted as the $z = 0$ incarnation of an equivalent twistor-type object. The twistor space is fibered over the complex z -line (or its compactification). The twistor space also has a $z = 1$ interpretation which is identified with the full solution of GLC.

NAHT and GLC

I want to explore the tantalizing possibility that the abelianized version of GLC is in fact equivalent, via recent breakthroughs in Non Abelian Hodge Theory (NAHT), to the full original (non-abelian) GLC, not only to its 0-th order or classical approximation.

Instead of viewing the solution constructed in [DP1] as a classical limit of the full solution, it is interpreted as the $z = 0$ incarnation of an equivalent twistor-type object. The twistor space is fibered over the complex z -line (or its compactification). The twistor space also has a $z = 1$ interpretation which is identified with the full solution of GLC.

This leads us to non-abelian Hodge theory and the connection with Simpson's talk.

NAHT

Non Abelian Hodge theory (NAHT), as developed by Hitchin, Donaldson, Corlette, Simpson, Saito, Sabbah, Mochizuki, and others, establishes under appropriate assumptions the equivalence of local systems and Higgs bundles.

NAHT

Non Abelian Hodge theory (NAHT), as developed by Hitchin, Donaldson, Corlette, Simpson, Saito, Sabbah, Mochizuki, and others, establishes under appropriate assumptions the equivalence of local systems and Higgs bundles. A richer object (harmonic bundle or twistor structure) is introduced, which specializes to both local systems and Higgs bundles.

NAHT

Non Abelian Hodge theory (NAHT), as developed by Hitchin, Donaldson, Corlette, Simpson, Saito, Sabbah, Mochizuki, and others, establishes under appropriate assumptions the equivalence of local systems and Higgs bundles. A richer object (harmonic bundle or twistor structure) is introduced, which specializes to both local systems and Higgs bundles. This is closely related to Deligne's notion of a λ -connection: at $\lambda = 1$ we have ordinary connections (or local systems), while at $\lambda = 0$ we have Higgs bundles. Depending on the exact context, these specialization maps are shown to be diffeomorphisms or categorical equivalences.

NAHT

The projective (or compact Kahler) case and the one dimensional open case were settled by Simpson twenty years ago - but the open case in higher dimension had to await the recent breakthroughs by Saito, Sabbah, Mochizuki, Jost-Yang-Zuo, Biquard, etc. This higher dimensional theory produces an equivalence of parabolic local systems and parabolic Higgs bundles. This is quite analogous to what is obtained in the compact case, except that the objects involved are required to satisfy three key conditions discovered by Mochizuki. Below we review these exciting developments, and outline our proposal for using NAHT to construct the automorphic sheaves required by the GLC.

NAHT, GLC, and QFT

This approach is purely mathematical of course, but it is parallel to physical ideas that have emerged from the collaborations of Witten with Kapustin, Gukov and Frenkel [KW, GW, W3, FW], where the GLC was placed firmly in the context of quantum field theory.

NAHT and GLC

Completion of these ideas depends on verification that Mochizuki's conditions are satisfied in situations arising from GLC. This requires a detailed analysis of instability loci in moduli spaces.

NAHT and GLC

Completion of these ideas depends on verification that Mochizuki's conditions are satisfied in situations arising from GLC. This requires a detailed analysis of instability loci in moduli spaces. Here we work with the moduli **spaces**, rather than the **stacks**. So stability is important, e.g. the difference between stability of a Higgs bundle and stability of the underlying bundle.

NAHT and GLC

Completion of these ideas depends on verification that Mochizuki's conditions are satisfied in situations arising from GLC. This requires a detailed analysis of instability loci in moduli spaces. Here we work with the moduli **spaces**, rather than the **stacks**. So stability is important, e.g. the difference between stability of a Higgs bundle and stability of the underlying bundle. Particularly important are the *Wobbly* locus of non-very-stable bundles, and the *Shaky* locus, roughly the Hitchin image of stable Higgs bundles with an unstable underlying bundle.

NAHT and GLC

Completion of these ideas depends on verification that Mochizuki's conditions are satisfied in situations arising from GLC. This requires a detailed analysis of instability loci in moduli spaces. Here we work with the moduli **spaces**, rather than the **stacks**. So stability is important, e.g. the difference between stability of a Higgs bundle and stability of the underlying bundle. Particularly important are the *Wobbly* locus of non-very-stable bundles, and the *Shaky* locus, roughly the Hitchin image of stable Higgs bundles with an unstable underlying bundle.

A bundle V is *very stable* if the only nilpotent Higgs field ϕ on V is $\phi = 0$.

NAHT and GLC

Completion of these ideas depends on verification that Mochizuki's conditions are satisfied in situations arising from GLC. This requires a detailed analysis of instability loci in moduli spaces. Here we work with the moduli **spaces**, rather than the **stacks**. So stability is important, e.g. the difference between stability of a Higgs bundle and stability of the underlying bundle. Particularly important are the *Wobbly* locus of non-very-stable bundles, and the *Shaky* locus, roughly the Hitchin image of stable Higgs bundles with an unstable underlying bundle.

A bundle V is *very stable* if the only nilpotent Higgs field ϕ on V is $\phi = 0$. (In other words, the cotangent fiber $T_{\{V\}}^* \mathbf{Bun}$ meets the Hitchin fiber over 0 only at the point $\phi = 0$.)

NAHT and GLC

Completion of these ideas depends on verification that Mochizuki's conditions are satisfied in situations arising from GLC. This requires a detailed analysis of instability loci in moduli spaces. Here we work with the moduli **spaces**, rather than the **stacks**. So stability is important, e.g. the difference between stability of a Higgs bundle and stability of the underlying bundle. Particularly important are the *Wobbly* locus of non-very-stable bundles, and the *Shaky* locus, roughly the Hitchin image of stable Higgs bundles with an unstable underlying bundle.

A bundle V is *very stable* if the only nilpotent Higgs field ϕ on V is $\phi = 0$. (In other words, the cotangent fiber $T_{\{V\}}^* \mathbf{Bun}$ meets the Hitchin fiber over 0 only at the point $\phi = 0$.) Laumon: very stable implies stable.

NAHT and GLC

Completion of these ideas depends on verification that Mochizuki's conditions are satisfied in situations arising from GLC. This requires a detailed analysis of instability loci in moduli spaces. Here we work with the moduli **spaces**, rather than the **stacks**. So stability is important, e.g. the difference between stability of a Higgs bundle and stability of the underlying bundle. Particularly important are the *Wobbly* locus of non-very-stable bundles, and the *Shaky* locus, roughly the Hitchin image of stable Higgs bundles with an unstable underlying bundle.

A bundle V is *very stable* if the only nilpotent Higgs field ϕ on V is $\phi = 0$. (In other words, the cotangent fiber $T_{\{V\}}^* \mathbf{Bun}$ meets the Hitchin fiber over 0 only at the point $\phi = 0$.) Laumon: very stable implies stable.

A bundle V is *wobbly* if it is stable but not very stable.

NAHT and GLC

Completion of these ideas depends on verification that Mochizuki's conditions are satisfied in situations arising from GLC. This requires a detailed analysis of instability loci in moduli spaces. Here we work with the moduli **spaces**, rather than the **stacks**. So stability is important, e.g. the difference between stability of a Higgs bundle and stability of the underlying bundle. Particularly important are the *Wobbly* locus of non-very-stable bundles, and the *Shaky* locus, roughly the Hitchin image of stable Higgs bundles with an unstable underlying bundle.

A bundle V is *very stable* if the only nilpotent Higgs field ϕ on V is $\phi = 0$. (In other words, the cotangent fiber $T_{\{V\}}^* \mathbf{Bun}$ meets the Hitchin fiber over 0 only at the point $\phi = 0$.) Laumon: very stable implies stable.

A bundle V is *wobbly* if it is stable but not very stable.

A bundle V is *shaky* if it is stable but there is a Higgs bundle (V', ϕ) mapping to it with V' unstable. (More details below.)

NAHT and GLC

Completion of these ideas depends on verification that Mochizuki's conditions are satisfied in situations arising from GLC. This requires a detailed analysis of instability loci in moduli spaces. Here we work with the moduli **spaces**, rather than the stacks. So stability is important, e.g. the difference between stability of a Higgs bundle and stability of the underlying bundle. Particularly important are the *Wobbly* locus of non-very-stable bundles, and the *Shaky* locus, roughly the Hitchin image of stable Higgs bundles with an unstable underlying bundle.

NAHT and GLC

Completion of these ideas depends on verification that Mochizuki's conditions are satisfied in situations arising from GLC. This requires a detailed analysis of instability loci in moduli spaces. Here we work with the moduli **spaces**, rather than the stacks. So stability is important, e.g. the difference between stability of a Higgs bundle and stability of the underlying bundle. Particularly important are the *Wobbly* locus of non-very-stable bundles, and the *Shaky* locus, roughly the Hitchin image of stable Higgs bundles with an unstable underlying bundle.

We will describe some known results about these loci for rank 2 bundles. These lead in some cases to an explicit construction (modulo solving the differential equations inherent in the non-abelian Hodge theory) of the Hecke eigensheaf demanded by the GLC.

NAHT: the theorems

Originally Corlette and Simpson proved the non-abelian Hodge theorem for projective manifolds:

Theorem [Corlette, Simpson] *Let $(X, \mathcal{O}_X(1))$ be a smooth complex projective variety. Then there is a natural equivalence of dg \otimes -categories:*

$$\mathbf{nah}_X : \left(\begin{array}{l} \text{finite rank } \mathbb{C}\text{-} \\ \text{local systems} \\ \text{on } X \end{array} \right) \longrightarrow \left(\begin{array}{l} \text{finite rank } \mathcal{O}_X(1)\text{-semistable} \\ \text{Higgs bundles on } X \text{ with} \\ ch_1 = 0 \text{ and } ch_2 = 0 \end{array} \right)$$

NAHT: the theorems

Mochizuki proved a version of the non-abelian Hodge correspondence which allows for singularities of the objects involved:

Theorem [Mochizuki] *Let $(X, \mathcal{O}_X(1))$ be a smooth complex projective variety and let $D \subset X$ be an effective divisor. Suppose that we have a closed subvariety $Z \subset X$ of codimension ≥ 3 , such that $X - Z$ is smooth and $D - Z$ is a normal crossing divisor. Then there is a canonical equivalence of dg \otimes -categories:*

$$\mathbf{nah}_{X,D} : \left(\begin{array}{l} \text{finite rank tame} \\ \text{parabolic } \mathbb{C}\text{-local} \\ \text{systems on } (X, D) \end{array} \right) \longrightarrow \left(\begin{array}{l} \text{finite rank locally abelian} \\ \text{tame parabolic Higgs} \\ \text{bundles on } (X, D) \\ \text{which are } \mathcal{O}_X(1)\text{-} \\ \text{semistable and satisfy} \\ \text{parch}_1 = 0 \text{ and } \text{parch}_2 = 0 \end{array} \right)$$

NAHT: the theorems

Mochizuki requires three basic ingredients for this theorem:

NAHT: the theorems

Mochizuki requires three basic ingredients for this theorem:

- (1) a good compactification, which is smooth and where the boundary is a divisor with normal crossings away from codimension 3;

NAHT: the theorems

Mochizuki requires three basic ingredients for this theorem:

- (1) a good compactification, which is smooth and where the boundary is a divisor with normal crossings away from codimension 3;
- (2) a local condition: tameness (the Higgs field is allowed to have at most logarithmic poles along D) and compatibility of filtrations (the parabolic structure is locally isomorphic to a direct sum of rank one objects); and

NAHT: the theorems

Mochizuki requires three basic ingredients for this theorem:

- (1) a good compactification, which is smooth and where the boundary is a divisor with normal crossings away from codimension 3;
- (2) a local condition: tameness (the Higgs field is allowed to have at most logarithmic poles along D) and compatibility of filtrations (the parabolic structure is locally isomorphic to a direct sum of rank one objects); and
- (3) a global condition: vanishing of parabolic Chern classes.

NAHT: the theorems

A feature of the non-abelian Hodge correspondence that is specific to the open case is captured in another result of Mochizuki:

Theorem [Mochizuki's Extension Theorem] *Let U be a quasi-projective variety and suppose U has two compactifications*

$$\phi : U \rightarrow X, \quad \psi : U \rightarrow Y$$

where:

- X, Y are projective and irreducible;
- X is smooth and $X - U$ is a normal crossing divisor away from codimension 3;

Then the restriction from X to U followed by the middle perversity extension from U to Y gives an equivalence of abelian categories:

$$\phi_{*!} \circ \psi^* : \left(\begin{array}{cc} \text{irreducible} & \text{tame} \\ \text{parabolic} & \mathbb{C}\text{-local} \\ \text{systems on } (X, D) & \end{array} \right) \longrightarrow \left(\begin{array}{l} \text{simple } \mathcal{D}\text{-modules on } Y \\ \text{which are smooth on } U \end{array} \right)$$

The plan

As we saw, non-abelian Hodge theory provides a natural approach to constructing the geometric Langlands correspondence \mathfrak{c} .

The plan

As we saw, non-abelian Hodge theory provides a natural approach to constructing the geometric Langlands correspondence \mathfrak{c} . The big hope is that the known eigensheaf of the abelianized Hecke, which is a Higgs-type object (\mathcal{E}, φ) , extends by non-abelian Hodge theory to a twistor eigensheaf on ${}^L\mathbf{Bun}$. The original Higgs sheaf appears at $z = 0$, while at the opposite end $z = 1$ we can expect to find precisely the Hecke eigensheaf postulated by the GLC.

The plan

The situation is essentially non-compact: In the moduli space ${}^L\mathbf{Bun}^S$ of stable bundles there is a locus \mathbf{S} of shaky bundles along which our Higgs field φ blows up. This can be traced back, essentially, to the difference between the notions of stability for bundles and Higgs bundles.

The plan

The situation is essentially non-compact: In the moduli space ${}^L\mathbf{Bun}^s$ of stable bundles there is a locus \mathbf{S} of shaky bundles along which our Higgs field φ blows up. This can be traced back, essentially, to the difference between the notions of stability for bundles and Higgs bundles. The cotangent bundle $T^\vee({}^L\mathbf{Bun}^s)$ embeds as a Zariski-open in ${}^L\mathbf{Higgs}^s$. If we ignore stability the two are equal: $T^\vee({}^L\mathbf{Bun}) = {}^L\mathbf{Higgs}$.

The plan

The situation is essentially non-compact: In the moduli space ${}^L\mathbf{Bun}^s$ of stable bundles there is a locus \mathbf{S} of shaky bundles along which our Higgs field φ blows up. This can be traced back, essentially, to the difference between the notions of stability for bundles and Higgs bundles. The cotangent bundle $T^\vee({}^L\mathbf{Bun}^s)$ embeds as a Zariski-open in ${}^L\mathbf{Higgs}^s$. If we ignore stability the two are equal: $T^\vee({}^L\mathbf{Bun}) = {}^L\mathbf{Higgs}$. But as moduli of stable objects, there is a locus \mathbf{Un} in ${}^L\mathbf{Higgs}$ parametrizing stable Higgs bundles with *unstable* underlying bundle. In order to turn the projection ${}^L\mathbf{Higgs}^s \rightarrow {}^L\mathbf{Bun}^s$ into a morphism, \mathbf{Un} must be blown up to an exceptional divisor $\widehat{\mathbf{Un}}$. Then the Higgs field part φ of the Hecke eigensheaf (\mathcal{E}, φ) on ${}^L\mathbf{Bun}^s$ blows up along the image \mathbf{S} of $\widehat{\mathbf{Un}}$.

The plan


So the heart of the matter amounts to verification of the Mochizuki conditions: we need to find where the Higgs field blows up, resolve this locus to obtain a normal crossing divisor, lift the objects to this resolution, and verify that the parabolic chern classes of these lifts vanish upstairs. This would provide the crucial third step in the following six step recipe for producing the candidate automorphic sheaf:

G -local system (V, ∇) on C

G -local system (V, ∇) on C

(1)

G -Higgs bundle (E, θ) on C



This is the Corlette-Simpson non-abelian Hodge correspondence $(E, \theta) = \mathbf{nah}_C(V, \nabla)$ on the smooth compact curve C .

G -local system (V, ∇) on C

(1)

G -Higgs bundle (E, θ) on C

G -local system (V, ∇) on C

(1)

G -Higgs bundle (E, θ) on C

(2)

${}_{ab}{}^L$ Hecke-eigensheaf on L Higgs

$$\boxed{G\text{-local system } (V, \nabla) \text{ on } C}$$

(1)

$$\boxed{G\text{-Higgs bundle } (E, \theta) \text{ on } C}$$

(2)

$$\boxed{{}_{ab}{}^L\text{Hecke-eigensheaf on } {}^L\text{Higgs}}$$

The functor (2) sends $(E, \theta) \in \mathbf{Higgs}$ to $\mathbf{FM}(\mathcal{O}_{(E, \theta)})$ where \mathbf{FM} is a Fourier-Mukai transform for coherent sheaves on $T^\vee \mathbf{Bun} = \mathbf{Higgs}$. In fact \mathbf{FM} is the integral transform with kernel the Poincaré sheaf constructed (away from the discriminant) in [DP1]. This sheaf is supported on the fiber product of the two Hitchin fibrations $h : \mathbf{Higgs}_0 \rightarrow B$ and ${}^L h : {}^L \mathbf{Higgs} \rightarrow B$.


G -local system (V, ∇) on C


(1)

G -Higgs bundle (E, θ) on C


(2)


${}_{ab}{}^L$ Hecke-eigensheaf on L Higgs







This part is known only in special cases, see below.










The functor (4) is the parabolic non-abelian Hodge correspondence $\mathbf{nah}_{{}^L \mathbf{Bun}^{ss}, S}$ of Mochizuki. Here ${}^L \mathbf{Bun}^{ss}$ denotes the (rigidified) stack of semistable bundles. Note that here we are applying the first Mochizuki theorem not to a projective variety but to a smooth proper Deligne-Mumford stack with a projective moduli space. In fact Mochizuki's proof works in this generality with no modifications.








Step 5 is just restriction.





The plan

Note that all of the other steps in this process are essentially already in place. The functor (1) is given by the Corlette-Simpson non-abelian Hodge correspondence $(E, \theta) = \mathbf{nah}_C(V, \nabla)$ on the smooth compact curve C .

The plan

Note that all of the other steps in this process are essentially already in place. The functor (1) is given by the Corlette-Simpson non-abelian Hodge correspondence $(E, \theta) = \mathbf{nah}_C(V, \nabla)$ on the smooth compact curve C .

The functor (2) sends $(E, \theta) \in \mathbf{Higgs}$ to $\mathbf{FM}(\mathcal{O}_{(E, \theta)})$ where \mathbf{FM} is a Fourier-Mukai transform for coherent sheaves on $T^\vee \mathbf{Bun} = \mathbf{Higgs}$. In fact \mathbf{FM} is the integral transform with kernel the Poincare sheaf constructed (away from the discriminant) in [DP1]. This sheaf is supported on the fiber product of the two Hitchin fibrations $h : \mathbf{Higgs}_0 \rightarrow B$ and ${}^L h : {}^L \mathbf{Higgs} \rightarrow B$.

The plan

Note that all of the other steps in this process are essentially already in place. The functor (1) is given by the Corlette-Simpson non-abelian Hodge correspondence $(E, \theta) = \mathbf{nah}_C(V, \nabla)$ on the smooth compact curve C .

The functor (2) sends $(E, \theta) \in \mathbf{Higgs}$ to $\mathbf{FM}(\mathcal{O}_{(E, \theta)})$ where \mathbf{FM} is a Fourier-Mukai transform for coherent sheaves on $T^\vee \mathbf{Bun} = \mathbf{Higgs}$. In fact \mathbf{FM} is the integral transform with kernel the Poincare sheaf constructed (away from the discriminant) in [DP1]. This sheaf is supported on the fiber product of the two Hitchin fibrations $h : \mathbf{Higgs}_0 \rightarrow B$ and ${}^L h : {}^L \mathbf{Higgs} \rightarrow B$.

The functor (4) is the parabolic non-abelian Hodge correspondence $\mathbf{nah}_{\mathbf{LBun}^{ss}, \mathbf{S}}$ of Mochizuki. Here ${}^L \mathbf{Bun}^{ss}$ denotes the (rigidified) stack of semistable bundles. Note that here we are applying the first Mochizuki theorem not to a projective variety but to a smooth proper Deligne-Mumford stack with a projective moduli space. In fact Mochizuki's proof works in this generality with no modifications.

The plan

The functors (5) and (6) are the pullback and middle extension functors applied to the two compactifications

${}^L\mathbf{Bun}^{ss} \supset {}^L\mathbf{Bun}^s \subset {}^L\mathbf{Bun}$. In order to conclude that the composition (6) \circ (5) is an equivalence we need a strengthening of Mochizuki's extension theorem which would allow for Y to be an Artin stack which is only locally of finite type.

Two forthcoming papers

In [DPS], with Pantev and Simpson, we study operations on Higgs bundles.

Two forthcoming papers

In [DPS], with Pantev and Simpson, we study operations on Higgs bundles. You have already heard about it, from Simpson.

Two forthcoming papers

In [DPS], with Pantev and Simpson, we study operations on Higgs bundles. You have already heard about it, from Simpson.

Natural operations such as pullback and direct image with respect to a morphism $f : X \rightarrow Y$ are well defined for harmonic bundles and local systems / D-modules, and commute with the NAHC. Pullback for Higgs bundles is also well defined and commutes with the NAHC.

The main goal is to give an algebraic definition of direct image for Higgs bundles and to show that it too commutes with the NAHC.

Two forthcoming papers

In [DPS], with Pantev and Simpson, we study operations on Higgs bundles. You have already heard about it, from Simpson.

Natural operations such as pullback and direct image with respect to a morphism $f : X \rightarrow Y$ are well defined for harmonic bundles and local systems / D-modules, and commute with the NAHC. Pullback for Higgs bundles is also well defined and commutes with the NAHC.

The main goal is to give an algebraic definition of direct image for Higgs bundles and to show that it too commutes with the NAHC.

The formula generalizes one of Zucker (for maps to a point). The proof uses Mochizuki's machinery, and a careful comparison of R-modules vs. parabolic modules.

Two forthcoming papers

In [DPS], with Pantev and Simpson, we study operations on Higgs bundles. You have already heard about it, from Simpson.

Natural operations such as pullback and direct image with respect to a morphism $f : X \rightarrow Y$ are well defined for harmonic bundles and local systems / D-modules, and commute with the NAHC. Pullback for Higgs bundles is also well defined and commutes with the NAHC.

The main goal is to give an algebraic definition of direct image for Higgs bundles and to show that it too commutes with the NAHC.

The formula generalizes one of Zucker (for maps to a point). The proof uses Mochizuki's machinery, and a careful comparison of R-modules vs. parabolic modules.

This formula allows us, in good situations, to compute the chern classes of parabolic Higgs sheaves on **Bun**. This is the missing step (3) in the flowchart. The results are then applied in [DP3] in a concrete example, namely P^1 with 5 parabolic points, to exhibit explicit solutions of GLC.


HAPPY BIRTHDAY, SAITO-SAN!!!



HAPPY BIRTHDAY, SAITO-SAN!!!



HAPPY BIRTHDAY, SAITO-SAN!!!



HAPPY BIRTHDAY, SAITO-SAN!!!



HAPPY BIRTHDAY, SAITO-SAN!!!



HAPPY BIRTHDAY, SAITO-SAN!!!



HAPPY BIRTHDAY, SAITO-SAN!!!



HAPPY BIRTHDAY, SAITO-SAN!!!

