

Discrete Hamiltonians of the discrete Painlevé equations

坂井 秀隆
SAKAI Hidetaka

Joint work with T. Mase and A. Nakamura

1 Oct. 2021

- 1 Introduction
- 2 discrete Painlevé equations
- 3 Case1: Biquadratic Hamiltonians of differential systems
- 4 Case2: $\#$ type
- 5 Case3: \sqsupset type

- 1 Introduction
- 2 discrete Painlevé equations
- 3 Case1: Biquadratic Hamiltonians of differential systems
- 4 Case2: $\#$ type
- 5 Case3: \sqsupset type

Introduction

We study the discrete Painlevé equations in parallel to studies of the Painlevé differential equations.

e.g. Lax pair, special solutions, and so on.

Introduction

We study the discrete Painlevé equations in parallel to studies of the Painlevé differential equations.

e.g. Lax pair, special solutions, and so on.

Difference:

We study the discrete Painlevé equations in parallel to studies of the Painlevé differential equations.

e.g. Lax pair, special solutions, and so on.

Difference:

The Painlevé differential equation is expressed as a Hamiltonian system, whereas the discrete Painlevé equation does not have such an expression.

The Painlevé differential equations are written in the form of the canonical systems with the Hamiltonians:

The Painlevé differential equations are written in the form of the canonical systems with the Hamiltonians:

$$\left[\frac{ds}{dt} = s(s-1) \right],$$

$$H_{VI} \left(\begin{matrix} a_1, a_2 \\ a_3, a_4 \end{matrix}; t; q, p \right) = q(q-1)(q-s)p^2 \\ + \left\{ (a_1 + 2a_2)q(q-1) + a_3(s-1)q + a_4s(q-1) \right\} p + a_2(a_1 + a_2)q,$$

$$H_V \left(\begin{matrix} a_1, a_2 \\ a_3 \end{matrix}; t; q, p \right) = p(p+1)q(q+e^t) + a_1q(p+1) + a_3pq - a_2e^tp,$$

$$H_{III}(D_6)(a_1, b_1; t; q, p) = p(p+1)q^2 - a_1p(q-1) - b_1pq - e^tq,$$

$$H_{III}(D_7)(a_1; t; q, p) = p^2q^2 + a_1qp + e^tp + q,$$

$$H_{III}(D_8)(t; q, p) = p^2q^2 + qp - q - \frac{e^t}{q},$$

$$H_{IV}(a_1, a_2; t; q, p) = pq(p-q-t) - a_1p - a_2q,$$

$$H_{II}(a_1; t; q, p) = p(p-q^2-t) - a_1q, \quad H_I(t; q, p) = p^2 - q^3 - tq.$$

Why do we use the Hamiltonian expression?

Why do we use the Hamiltonian expression?

1. Hamiltonian is an invariant.

Why do we use the Hamiltonian expression?

1. Hamiltonian is an invariant.

→ It is not applicable to non-autonomous cases.

Why do we use the Hamiltonian expression?

1. Hamiltonian is an invariant.
→ It is not applicable to non-autonomous cases.
2. Hamiltonian gives a simple expression to a dynamical system.

Why do we use the Hamiltonian expression?

1. Hamiltonian is an invariant.
→ It is not applicable to non-autonomous cases.
2. Hamiltonian gives a simple expression to a dynamical system.
→ The Hamiltonian function can give a clue to identify the system.

Why do we use the Hamiltonian expression?

1. Hamiltonian is an invariant.
→ It is not applicable to non-autonomous cases.
2. Hamiltonian gives a simple expression to a dynamical system.
→ The Hamiltonian function can give a clue to identify the system.

If a discrete dynamical system can be described simply using function W on a phase space, we call this W a discrete Hamiltonian, although it is a vague terminology.

Discrete Lagrangian and discrete Hamiltonian

Starting from a function: $L_k(r, s) : M^n \times M^n \rightarrow \mathbb{R}$. (We call it Lagrangian.)

Discrete Lagrangian and discrete Hamiltonian

Starting from a function: $L_k(r, s) : M^n \times M^n \rightarrow \mathbb{R}$. (We call it Lagrangian.)

We consider formal sum $S(\lambda) = \sum_{k \in \mathbb{Z}} L_k(\lambda_k, \lambda_{k+1})$, and $\delta S = 0$:

$$\begin{aligned}\delta S(\lambda) &= \sum_{k \in \mathbb{Z}} \delta L_k(\lambda_k, \lambda_{k+1}) \\&= \sum_{k \in \mathbb{Z}} \{L_k(\lambda_k + \delta \lambda_k, \lambda_{k+1} + \delta \lambda_{k+1}) - L_k(\lambda_k, \lambda_{k+1})\} \\&= \sum_{k \in \mathbb{Z}} \left\{ \frac{\partial L_k}{\partial r}(\lambda_k, \lambda_{k+1}) \delta \lambda_k + \frac{\partial L_k}{\partial s}(\lambda_k, \lambda_{k+1}) \delta \lambda_{k+1} \right\} \\&= \sum_{k \in \mathbb{Z}} \left\{ \frac{\partial L_k}{\partial r}(\lambda_k, \lambda_{k+1}) + \frac{\partial L_{k-1}}{\partial s}(\lambda_{k-1}, \lambda_k) \right\} \delta \lambda_k = 0.\end{aligned}$$

Discrete Lagrangian and discrete Hamiltonian

Starting from a function: $L_k(r, s) : M^n \times M^n \rightarrow \mathbb{R}$. (We call it Lagrangian.)

We consider formal sum $S(\lambda) = \sum_{k \in \mathbb{Z}} L_k(\lambda_k, \lambda_{k+1})$, and $\delta S = 0$:

$$\begin{aligned}\delta S(\lambda) &= \sum_{k \in \mathbb{Z}} \delta L_k(\lambda_k, \lambda_{k+1}) \\&= \sum_{k \in \mathbb{Z}} \{L_k(\lambda_k + \delta \lambda_k, \lambda_{k+1} + \delta \lambda_{k+1}) - L_k(\lambda_k, \lambda_{k+1})\} \\&= \sum_{k \in \mathbb{Z}} \left\{ \frac{\partial L_k}{\partial r}(\lambda_k, \lambda_{k+1}) \delta \lambda_k + \frac{\partial L_k}{\partial s}(\lambda_k, \lambda_{k+1}) \delta \lambda_{k+1} \right\} \\&= \sum_{k \in \mathbb{Z}} \left\{ \frac{\partial L_k}{\partial r}(\lambda_k, \lambda_{k+1}) + \frac{\partial L_{k-1}}{\partial s}(\lambda_{k-1}, \lambda_k) \right\} \delta \lambda_k = 0.\end{aligned}$$

$$\rightarrow \quad \frac{\partial L_k}{\partial r}(\lambda_k, \lambda_{k+1}) + \frac{\partial L_{k-1}}{\partial s}(\lambda_{k-1}, \lambda_k) = 0 \quad (\text{Euler-Lagrange}).$$

Discrete Lagrangian and discrete Hamiltonian

Legendre transformation:

We put $\mu_k = \frac{\partial L_k}{\partial r}(\lambda_k, \lambda_{k+1}) = -\frac{\partial L_{k-1}}{\partial s}(\lambda_{k-1}, \lambda_k)$,
and put $H(\lambda, \bar{\mu}) = \bar{\lambda}\bar{\mu} + L(\lambda, \bar{\lambda})$.

Discrete Lagrangian and discrete Hamiltonian

Legendre transformation:

We put $\mu_k = \frac{\partial L_k}{\partial r}(\lambda_k, \lambda_{k+1}) = -\frac{\partial L_{k-1}}{\partial s}(\lambda_{k-1}, \lambda_k)$,
and put $H(\lambda, \bar{\mu}) = \bar{\lambda}\bar{\mu} + L(\lambda, \bar{\lambda})$.

Then the system is written as

$$\bar{\lambda} = \frac{\partial H}{\partial \bar{\mu}}, \quad \mu = \frac{\partial H}{\partial \lambda}.$$

Discrete Lagrangian and discrete Hamiltonian

Legendre transformation:

We put $\mu_k = \frac{\partial L_k}{\partial r}(\lambda_k, \lambda_{k+1}) = -\frac{\partial L_{k-1}}{\partial s}(\lambda_{k-1}, \lambda_k)$,
and put $H(\lambda, \bar{\mu}) = \bar{\lambda}\bar{\mu} + L(\lambda, \bar{\lambda})$.

Then the system is written as

$$\bar{\lambda} = \frac{\partial H}{\partial \bar{\mu}}, \quad \mu = \frac{\partial H}{\partial \lambda}.$$

But we know this form. It is just a generating function of a canonical transformation.

$d\mu \wedge d\lambda$ is an invariant form (symplectic).

Discrete Lagrangian and discrete Hamiltonian

Legendre transformation:

We put $\mu_k = \frac{\partial L_k}{\partial r}(\lambda_k, \lambda_{k+1}) = -\frac{\partial L_{k-1}}{\partial s}(\lambda_{k-1}, \lambda_k)$,
and put $H(\lambda, \bar{\mu}) = \bar{\lambda}\bar{\mu} + L(\lambda, \bar{\lambda})$.

Then the system is written as

$$\bar{\lambda} = \frac{\partial H}{\partial \bar{\mu}}, \quad \mu = \frac{\partial H}{\partial \lambda}.$$

But we know this form. It is just a generating function of a canonical transformation.

$d\mu \wedge d\lambda$ is an invariant form (symplectic).

We forget the Lagrangian, and only consider the generating function of the canonical transformation.

- 1 Introduction
- 2 discrete Painlevé equations
- 3 Case1: Biquadratic Hamiltonians of differential systems
- 4 Case2: $\#$ type
- 5 Case3: \sqsupset type

Discrete Painlevé equations are classified by the types of certain rational surfaces:

Discrete Painlevé equations are classified by the types of certain rational surfaces:

Definition

Let X be a smooth projective rational surface. We call X a generalized Halphen surface if X has an anti-canonical divisor of canonical type.

discrete Painlevé equations

Discrete Painlevé equations are classified by the types of certain rational surfaces:

Definition

Let X be a smooth projective rational surface. We call X a generalized Halphen surface if X has an anti-canonical divisor of canonical type.

Let $D = \sum_{i \in I} m_i D_i$ be an effective divisor on X with irreducible components D_i . We say that D is of canonical type if

$$\mathcal{K}_X \cdot [D_i] = 0 \quad \text{for all } i.$$

- $\dim |-\mathcal{K}_X| = 1 \rightarrow$ rational elliptic surface
- $\dim |-\mathcal{K}_X| = 0$

- $\dim |-\mathcal{K}_X| = 1 \rightarrow$ rational elliptic surface
- $\dim |-\mathcal{K}_X| = 0 \rightarrow$ Painlevé

- $\dim |-\mathcal{K}_X| = 1 \rightarrow$ rational elliptic surface
- $\dim |-\mathcal{K}_X| = 0 \rightarrow$ Painlevé

Classification by anti-canonical divisor

elliptic	$A_0^{(1)}$
multiplicative	$A_0^{(1)*}, A_1^{(1)}, A_2^{(1)},$ $A_3^{(1)}, \dots, A_6^{(1)}, A_7^{(1)}, A_7^{(1)'}, A_8^{(1)}$
additive	$A_0^{(1)**}, A_1^{(1)*}, A_2^{(1)*},$ $D_4^{(1)}, D_5^{(1)}, D_6^{(1)}, D_7^{(1)}, D_8^{(1)},$ $E_6^{(1)}, E_7^{(1)}, E_8^{(1)}$

We have non-autonomous differential systems only for $D_k^{(1)}$ and $E_k^{(1)}$.

equations	P_{VI}	P_V	$P_{III}(D_6)$	$P_{III}(D_7)$	$P_{III}(D_8)$
geometry	$D_4^{(1)}$	$D_5^{(1)}$	$D_6^{(1)}$	$D_7^{(1)}$	$D_8^{(1)}$
symmetry	$D_4^{(1)}$	$A_3^{(1)}$	$(A_1 + A_1)^{(1)}$	$A_1^{(1)}$	-

P_{IV}	P_{II}	P_I
$E_6^{(1)}$	$E_7^{(1)}$	$E_8^{(1)}$
$A_2^{(1)}$	$A_1^{(1)}$	-

Definition

Discrete Painlevé equation is a discrete dynamical system which is given by a Cremona isometry of a generalized Halphen surface of infinite order.

Definition

Discrete Painlevé equation is a discrete dynamical system which is given by a Cremona isometry of a generalized Halphen surface of infinite order.

We have discrete Painlevé equations for each surfaces except $A_8^{(1)}$, $D_8^{(1)}$, $E_8^{(1)}$.

Definition

Discrete Painlevé equation is a discrete dynamical system which is given by a Cremona isometry of a generalized Halphen surface of infinite order.

We have discrete Painlevé equations for each surfaces except $A_8^{(1)}$, $D_8^{(1)}$, $E_8^{(1)}$.

Except $E_8^{(1)}$, the surfaces can be blown down to $\mathbb{P}^1 \times \mathbb{P}^1$. We set the coordinate as $(f_0 : f_1), (g_0 : g_1)$.

We divide them into 4 cases:

- 1 The image of the anti-canonical divisor is $f_0^2 g_0^2 = 0$,
- 2 The image of the anti-canonical divisor is $f_0 f_1 g_0^2 = 0$,
- 3 The image of the anti-canonical divisor is $f_0 f_1 g_0 g_1 = 0$,
- 4 The others.

- ① $f_0^2 g_0^2 = 0$: $D_5^{(1)}, D_6^{(1)}, D_7^{(1)}, E_6^{(1)}, E_7^{(1)},$
- ② $f_0 f_1 g_0^2 = 0$: $D_4^{(1)}, D_5^{(1)}, D_6^{(1)}, D_7^{(1)}, (D_8^{(1)}),$
- ③ $f_0 f_1 g_0 g_1 = 0$: $A_3^{(1)}, A_4^{(1)}, A_5^{(1)}, A_6^{(1)}, A_7^{(1)}, A_7^{(1)'}, (A_8^{(1)}),$
- ④ the others: $A_0^{(1)}, A_0^{(1)*}, A_0^{(1)**}, A_1^{(1)}, A_1^{(1)*}, A_2^{(1)}, A_2^{(1)*}.$

- 1 Introduction
- 2 discrete Painlevé equations
- 3 Case1: Biquadratic Hamiltonians of differential systems**
- 4 Case2: $\#$ type
- 5 Case3: \supset type

Case1: $f_0^2 g_0^2 = 0$

In this case, the Hamiltonian of the differential system is written by a biquadratic polynomial:

$$H = (g^2, g, 1) \begin{pmatrix} m_{22} & m_{21} & m_{20} \\ m_{12} & m_{11} & m_{10} \\ m_{02} & m_{01} & m_{00} \end{pmatrix} \begin{pmatrix} f^2 \\ f \\ 1 \end{pmatrix},$$
$$\frac{df}{dt} = \frac{\partial H}{\partial g}, \quad \frac{dg}{dt} = -\frac{\partial H}{\partial f}.$$

$$M = M_{D_5} = \begin{pmatrix} 1 & s & 0 \\ 1 & s + a_1 + a_3 & -a_2 s \\ 0 & a_1 & 0 \end{pmatrix}, \quad M_{D_6} = \begin{pmatrix} 1 & 0 & 0 \\ 1 & -a_1 - b_1 & -s \\ 0 & -a_1 & 0 \end{pmatrix},$$

$$M_{D_7} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & a_1 & s \\ 0 & 1 & 0 \end{pmatrix}, \quad M_{E_6} = \begin{pmatrix} 0 & 1 & 0 \\ -1 & -s & -a_2 \\ 0 & -a_1 & 0 \end{pmatrix}, \quad M_{E_7} = \begin{pmatrix} 0 & 0 & 1 \\ -1 & 0 & -s \\ 0 & -a_1 & 0 \end{pmatrix}.$$

$s = e^t$ for D -type, $s = t$ for E -type.

Discrete Painlevé equations are written by using these matrices as:

$$g = -\bar{g} - \frac{\hat{m}_{12}f^2 + \hat{m}_{11}f + \hat{m}_{10}}{\hat{m}_{22}f^2 + \hat{m}_{21}f + \hat{m}_{20}}, \quad \bar{f} = -f - \frac{\bar{m}_{21}\bar{g}^2 + \bar{m}_{11}\bar{g} + \bar{m}_{01}}{\bar{m}_{22}\bar{g}^2 + \bar{m}_{12}\bar{g} + \bar{m}_{02}}.$$

Discrete Painlevé equations are written by using these matrices as:

$$g = -\bar{g} - \frac{\hat{m}_{12}f^2 + \hat{m}_{11}f + \hat{m}_{10}}{\hat{m}_{22}f^2 + \hat{m}_{21}f + \hat{m}_{20}}, \quad \bar{f} = -f - \frac{\bar{m}_{21}\bar{g}^2 + \bar{m}_{11}\bar{g} + \bar{m}_{01}}{\bar{m}_{22}\bar{g}^2 + \bar{m}_{12}\bar{g} + \bar{m}_{02}}.$$

Hence when we put a generating function W as

$$W(f, \bar{g}) = -f\bar{g} - \int \frac{\hat{m}_{12}f^2 + \hat{m}_{11}f + \hat{m}_{10}}{\hat{m}_{22}f^2 + \hat{m}_{21}f + \hat{m}_{20}} df - \int \frac{\bar{m}_{21}\bar{g}^2 + \bar{m}_{11}\bar{g} + \bar{m}_{01}}{\bar{m}_{22}\bar{g}^2 + \bar{m}_{12}\bar{g} + \bar{m}_{02}} d\bar{g},$$

we can write the discrete equations as

$$g = \frac{\partial W}{\partial f}, \quad \bar{f} = \frac{\partial W}{\partial \bar{g}}.$$

The explicit formula:

$$W = W_{D_5} = -f\bar{g} - f - s\bar{g} - \bar{a}_3 \log(\bar{g} + 1) + a_2 \log f \\ - \bar{a}_1 \log \bar{g} - (a_1 + a_2 + a_3 - 1) \log(f + s),$$

$$W_{D_6} = -f\bar{g} - f - \frac{s}{f} + (a_1 + b_1 - 1) \log f \\ + \bar{a}_1 \log \bar{g} + \bar{b}_1 \log(\bar{g} + 1),$$

$$W_{D_7} = -f\bar{g} - \frac{s}{f} + \frac{1}{\bar{g}} - (a_1 - 1) \log f - \bar{a}_1 \log \bar{g},$$

$$W_{E_6} = -f\bar{g} + \frac{f^2}{2} + sf + \frac{\bar{g}^2}{2} - s\bar{g} + a_2 \log f - \bar{a}_1 \log \bar{g},$$

$$W_{E_7} = -f\bar{g} + sf + \frac{f^3}{3} - \bar{a}_1 \log \bar{g}.$$

e.g. $E_7^{(1)}$ type:

$$W_{E_7} = -f\bar{g} + sf + \frac{f^3}{3} - \bar{a}_1 \log \bar{g},$$

e.g. $E_7^{(1)}$ type:

$$W_{E_7} = -f\bar{g} + sf + \frac{f^3}{3} - \bar{a}_1 \log \bar{g},$$

$$g = \frac{\partial W_{E_7}}{\partial f} = -\bar{g} + s + f^2,$$

$$\bar{f} = \frac{\partial W_{E_7}}{\partial \bar{g}} = -f - \frac{\bar{a}_1}{\bar{g}}.$$

Content

- 1 Introduction
- 2 discrete Painlevé equations
- 3 Case1: Biquadratic Hamiltonians of differential systems
- 4 Case2: $\#$ type**
- 5 Case3: \sqsupset type

Case2: $f_0 f_1 g_0 g_1 = 0$

$$M_{A_7} = \begin{pmatrix} 0 & -a_0 & 0 \\ 1 & 0 & 0 \\ 0 & -1 & 1 \end{pmatrix}, \quad M_{A'_7} = \begin{pmatrix} 1 & -a_0 & 0 \\ 0 & 0 & 0 \\ 0 & -1 & 1 \end{pmatrix},$$

$$M_{A_6} = \begin{pmatrix} 0 & 1/b & 0 \\ 1 & 0 & -1/b \\ 0 & -a_1 & a_1 \end{pmatrix}, \quad M_{A_5} = \begin{pmatrix} 0 & b_1/a_2 & 0 \\ a_0 & 0 & -b_1/a_2 \\ 1/a_1 & -1 - (1/a_1) & 1 \end{pmatrix},$$

$$M_{A_4} = \begin{pmatrix} 0 & 1 & -1 \\ a_0/a_2 & 0 & 1 + (1/a_4) \\ -a_0 a_3/a_2 & a_0 a_3 + (1/a_2 a_4) & -1/a_4 \end{pmatrix},$$

$$M_{A_3} = \begin{pmatrix} a_0 a_5 & -1/(a_1 a_2^2 a_3) - a_0 a_3 a_5 & 1/(a_1 a_2^2) \\ -(1 + a_0) a_5 & 0 & -(1 + a_1)/a_1 a_2 \\ a_5 & -1 - a_5 & 1 \end{pmatrix}.$$

Discrete Painlevé equations are written by using these matrices as:

$$g = \frac{\hat{m}_{02}f^2 + \hat{m}_{01}f + \hat{m}_{00}}{\bar{g}(\hat{m}_{22}f^2 + \hat{m}_{21}f + \hat{m}_{20})}, \quad \bar{f} = \frac{\bar{m}_{20}\bar{g}^2 + \bar{m}_{10}\bar{g} + \bar{m}_{00}}{f(\bar{m}_{22}\bar{g}^2 + \bar{m}_{12}\bar{g} + \bar{m}_{02})}$$

Discrete Painlevé equations are written by using these matrices as:

$$g = \frac{\hat{m}_{02}f^2 + \hat{m}_{01}f + \hat{m}_{00}}{\bar{g}(\hat{m}_{22}f^2 + \hat{m}_{21}f + \hat{m}_{20})}, \quad \bar{f} = \frac{\bar{m}_{20}\bar{g}^2 + \bar{m}_{10}\bar{g} + \bar{m}_{00}}{f(\bar{m}_{22}\bar{g}^2 + \bar{m}_{12}\bar{g} + \bar{m}_{02})}$$

The symplectic form is $\omega = \frac{dg \wedge df}{fg} = d \log g \wedge d \log f$. When we put $F = \log f$, $G = \log g$, we can find a generating function $\widetilde{W}(F, \bar{G})$. But it is important that the system is a birational mapping, so we want to use the variables f and g .

We put $W(f, \bar{g}) = \widetilde{W}(\log f, \log \bar{g})$, then

$$\begin{aligned} W(f, \bar{g}) = & -\log f \log \bar{g} + \int \log (\hat{m}_{02} f^2 + \hat{m}_{01} f + \hat{m}_{00}) \frac{df}{f} \\ & - \int \log (\hat{m}_{22} f^2 + \hat{m}_{21} f + \hat{m}_{20}) \frac{df}{f} \\ & + \int \log (\bar{m}_{20} \bar{g}^2 + \bar{m}_{10} \bar{g} + \bar{m}_{00}) \frac{d\bar{g}}{\bar{g}} \\ & - \int \log (\bar{m}_{22} \bar{g}^2 + \bar{m}_{12} \bar{g} + \bar{m}_{02}) \frac{d\bar{g}}{\bar{g}}, \end{aligned}$$

and we can write the discrete equations as

$$g = \exp \left(f \frac{\partial W}{\partial f} \right), \quad \bar{f} = \exp \left(\bar{g} \frac{\partial W}{\partial \bar{g}} \right).$$

The explicit formula:

$$\begin{aligned}
 W_{A_3} = & -\log f \log \bar{g} + \operatorname{Li}_2(\bar{g}) + \operatorname{Li}_2(\bar{a}_0 \bar{g}) - \operatorname{Li}_2\left(\frac{\bar{g}}{\bar{a}_2}\right) - \operatorname{Li}_2\left(\frac{\bar{g}}{\bar{a}_1 \bar{a}_2}\right) \\
 & - \operatorname{Li}_2(f) + \operatorname{Li}_2\left(\frac{f}{a_3}\right) - \operatorname{Li}_2(a_5 f) + \operatorname{Li}_2\left(\frac{a_0 a_1 a_2^2 a_3 a_5 f}{q}\right) \\
 & - \log \bar{a}_3 \log \bar{g} + \log(a_1 a_2^2) \log f,
 \end{aligned}$$

$$\begin{aligned}
 W_{A_4} = & -\log f \log \bar{g} + \operatorname{Li}_2(f) - \operatorname{Li}_2\left(\frac{qf}{a_2}\right) - \operatorname{Li}_2(a_0 a_3 a_4 f) - \operatorname{Li}_2(\bar{g}) \\
 & - \operatorname{Li}_2(\bar{a}_4 \bar{g}) + \operatorname{Li}_2\left(\frac{\bar{g}}{\bar{a}_3}\right) + \left(\log \frac{\bar{a}_2}{\bar{a}_0 \bar{a}_3 \bar{a}_4}\right) \log \bar{g} - \log a_4 \log f,
 \end{aligned}$$

$$\begin{aligned}
 W_{A_5} = & -\log f \log \bar{g} - \operatorname{Li}_2(f) - \operatorname{Li}_2\left(\frac{f}{a_1}\right) - \operatorname{Li}_2\left(\frac{\bar{b}_1 \bar{g}}{\bar{a}_2}\right) \\
 & + \operatorname{Li}_2(-\bar{a}_0 \bar{a}_1 \bar{g}) - \frac{1}{2} \left(\log \frac{b_1 f}{a_1}\right)^2 + \log \bar{a}_1 \log \bar{g},
 \end{aligned}$$

$$\begin{aligned}
W_{A_6} &= -\log f \log \bar{g} - \text{Li}_2(f) - \text{Li}_2\left(\frac{\bar{g}}{\bar{a}_1 \bar{b}}\right) + \log \bar{g} \log \bar{a}_1 \\
&\quad - \frac{1}{2} \left(\log \frac{f}{qb} \right)^2 - \frac{1}{2} (\log \bar{g})^2 - \log a_1 \log f, \\
W_{A'_7} &= -\log f \log \bar{g} - \frac{1}{2} (\log f)^2 - (\log \bar{g})^2 - \text{Li}_2(f) + \text{Li}_2\left(\frac{qf}{a_0}\right) \\
&\quad - \log \frac{-a_0}{q} \log f, \\
W_{A_7} &= -\log f \log \bar{g} - \text{Li}_2(f) - \frac{1}{2} (\log(-q^{-1} a_0 f))^2 - \frac{1}{2} (\log(\bar{g}))^2,
\end{aligned}$$

where $\text{Li}_2(x)$ is the dilogarithmic function:

$$\text{Li}_2(x) = - \int \frac{\log(1-x)}{x} dx = \sum_{k=1}^{\infty} \frac{x^k}{k^n}.$$

e.g. $A_7^{(1) '}$ type:

$$W_{A_7'} = -\log f \log \bar{g} - \frac{1}{2}(\log f)^2 - (\log \bar{g})^2 - \text{Li}_2(f) + \text{Li}_2\left(\frac{qf}{a_0}\right) - \log \frac{-a_0}{q} \log f,$$

e.g. $A_7^{(1)'} \text{ type:}$

$$W_{A'_7} = -\log f \log \bar{g} - \frac{1}{2}(\log f)^2 - (\log \bar{g})^2 - \text{Li}_2(f) + \text{Li}_2\left(\frac{qf}{a_0}\right) - \log \frac{-a_0}{q} \log f,$$

$$\begin{aligned} g &= \exp\left(f \frac{\partial W_{A'_7}}{\partial f}\right) \\ &= \exp\left(-\log \bar{g} - \log f + \log(1-f) - \log\left(1 - \frac{qf}{a_0}\right) - \log \frac{-a_0}{q}\right) \\ &= \frac{1-f}{\bar{g}f\left(f - \frac{a_0}{q}\right)}, \\ \bar{f} &= \exp\left(\bar{g} \frac{\partial W_{A'_7}}{\partial \bar{g}}\right) = \exp(-\log f - 2\log \bar{g}) = \frac{1}{f\bar{g}^2}. \end{aligned}$$

Content

- 1 Introduction
- 2 discrete Painlevé equations
- 3 Case1: Biquadratic Hamiltonians of differential systems
- 4 Case2: $\#$ type
- 5 Case3: \sqsupset type

Case3: $f_0 f_1 g_0^2 = 0$

$$M_{D_4} = \begin{pmatrix} 1 & -1-s & s \\ a_1 + 2a_2 & -a_1 - 2a_2 + (s-1)a_3 + a_4 & sa_4 \\ a_2(a_1 + a_2) & 0 & 0 \end{pmatrix}.$$

Discrete Painlevé equation is written by using the matrix as:

$$g = -\bar{g} - \frac{m_{12}f^2 + m_{11}f + m_{10}}{m_{22}f^2 + m_{21}f + m_{20}}, \quad \bar{f} = \frac{\bar{m}_{20}\bar{g}^2 + \bar{m}_{10}\bar{g} + \bar{m}_{00}}{f(\bar{m}_{22}\bar{g}^2 + \bar{m}_{12}\bar{g} + \bar{m}_{02})}.$$

Case3: $f_0 f_1 g_0^2 = 0$

$$M_{D_4} = \begin{pmatrix} 1 & -1-s & s \\ a_1 + 2a_2 & -a_1 - 2a_2 + (s-1)a_3 + a_4 & sa_4 \\ a_2(a_1 + a_2) & 0 & 0 \end{pmatrix}.$$

Discrete Painlevé equation is written by using the matrix as:

$$g = -\bar{g} - \frac{m_{12}f^2 + m_{11}f + m_{10}}{m_{22}f^2 + m_{21}f + m_{20}}, \quad \bar{f} = \frac{\bar{m}_{20}\bar{g}^2 + \bar{m}_{10}\bar{g} + \bar{m}_{00}}{f(\bar{m}_{22}\bar{g}^2 + \bar{m}_{12}\bar{g} + \bar{m}_{02})}.$$

Hence when we put a function W as

$$\begin{aligned} W(f, \bar{g}) = & -\bar{g} \log f - \int \frac{m_{12}f^2 + m_{11}f + m_{10}}{m_{22}f^2 + m_{21}f + m_{20}} \frac{df}{f} \\ & + \int \log (\bar{m}_{20}\bar{g}^2 + \bar{m}_{10}\bar{g} + \bar{m}_{00}) d\bar{g} \\ & - \int \log (\bar{m}_{22}\bar{g}^2 + \bar{m}_{12}\bar{g} + \bar{m}_{02}) d\bar{g}, \end{aligned}$$

we can write the discrete equations as

$$g = f \frac{\partial W}{\partial f}, \quad \bar{f} = \exp \left(\frac{\partial W}{\partial \bar{g}} \right).$$

e.g. $D_4^{(1) '}$ type:

$$\begin{aligned} W_{D_4} = & -\bar{g} \log f + a_4 \log f - a_3 \log(1 - f) \\ & - (a_1 + 2a_2 + a_3 + a_4 - 1) \log(1 - f/s) + \bar{g}(\log \bar{g} + \log s) \\ & - (\bar{g} + \bar{a}_1 + \bar{a}_2) \log(\bar{g} + \bar{a}_1 + \bar{a}_2) - (\bar{g} + \bar{a}_2) \log(\bar{g} + \bar{a}_2) \\ & + (\bar{g} - \bar{a}_4) \log(\bar{g} - \bar{a}_4), \end{aligned}$$

e.g. $D_4^{(1)'} type:$

$$\begin{aligned}
 W_{D_4} = & -\bar{g} \log f + a_4 \log f - a_3 \log(1-f) \\
 & - (a_1 + 2a_2 + a_3 + a_4 - 1) \log(1-f/s) + \bar{g}(\log \bar{g} + \log s) \\
 & - (\bar{g} + \bar{a}_1 + \bar{a}_2) \log(\bar{g} + \bar{a}_1 + \bar{a}_2) - (\bar{g} + \bar{a}_2) \log(\bar{g} + \bar{a}_2) \\
 & + (\bar{g} - \bar{a}_4) \log(\bar{g} - \bar{a}_4),
 \end{aligned}$$

$$\begin{aligned}
 g = f \frac{\partial W_{D_4}}{\partial f} &= -\bar{g} + a_4 - \frac{a_3 f}{1-f} + \frac{(a_1 + 2a_2 + a_3 + a_4 - 1)f}{s-f} \\
 &= -\bar{g} + 1 - a_1 - 2a_2 - \frac{a_3}{1-f} + \frac{a_1 + 2a_2 + a_3 + a_4 - 1}{1-f/s}
 \end{aligned}$$

$$\begin{aligned}
 \bar{f} &= \exp \left(\frac{\partial W_{D_4}}{\partial \bar{g}} \right) \\
 &= \exp \left(-\log f + \log \bar{g} + \log(\bar{g} - \bar{a}_4) + \log s - \log(\bar{g} + \bar{a}_1 + \bar{a}_2) \right. \\
 &\quad \left. - \log(\bar{g} + \bar{a}_2) \right) \\
 &= \frac{s \bar{g} (\bar{g} - \bar{a}_4)}{f (\bar{g} + \bar{a}_1 + \bar{a}_2) (\bar{g} + \bar{a}_2)},
 \end{aligned}$$

Thank you.