

# Gauge-Higgs Unification

H parity and LHC/ILC signals



Yutaka Hosotani, Osaka University

Workshop at Kobe University, 6 January 2011

# Gauge-Higgs Unification in 5 dimensions

4-dim. components  $A_\mu$

extra-dim. component  $A_y$

Hosotani 1983, 1989  
Davies, McLachlan 1988, 1989  
Hatanaka, Inami, Lim, 1998

AB phase  
in extra dim

$$e^{i\hat{\theta}_H(x)} \sim P \exp \left\{ ig \int_C dy A_y \right\}$$

$$\hat{\theta}_H(x) = \theta_H + \frac{H(x)}{f_H} \quad \text{Higgs field} = \text{gauge field}$$

*stable Higgs*

*precision  
measurements*


*KK modes*

# $SO(5) \times U(1)$ in Randall-Sundrum warped space

$$ds^2 = e^{-2k|y|} dx_\mu dx^\mu + dy^2$$

$$0 \leq |y| \leq L = \pi R$$

Agashe, Contino, Pomarol 2005  
 Hosotani, Sakamura 2006  
 Medina, Shah, Wagner 2007



*Orbifold BC*

$$\begin{pmatrix} A_\mu \\ A_y \end{pmatrix} (x, -y) = P_0 \begin{pmatrix} A_\mu \\ -A_y \end{pmatrix} (x, y) P_0^\dagger$$

$$\begin{pmatrix} A_\mu \\ A_y \end{pmatrix} (x, \pi R - y) = P_1 \begin{pmatrix} A_\mu \\ -A_y \end{pmatrix} (x, \pi R + y) P_1^\dagger$$

# 4D gauge bosons and Higgs

$$P_0 = P_1 = \begin{pmatrix} -1 & & & & \\ & -1 & & & \\ & & -1 & & \\ & & & -1 & \\ & & & & +1 \end{pmatrix}$$



*W Z γ*

$A_\mu \sim$



$$SO(5) \rightarrow SO(4) \simeq SU(2)_L \times SU(2)_R$$



*Higgs*

$A_y \sim$

$$\begin{pmatrix} \begin{matrix} \phi_1 \\ \phi_2 \\ \phi_3 \\ \phi_4 \end{matrix} \\ \square \end{pmatrix}$$

$$\Phi = \begin{bmatrix} \phi_1 + i\phi_2 \\ \phi_4 - i\phi_3 \end{bmatrix}$$

4D Higgs doublet

# Matter content

YH, Noda, Uekusa 2009

(YH, Oda, Ohnuma, Sakamura 2008)

Planck brane

$SO(5) \times U(1)$

TeV brane

Quarks

$$\begin{pmatrix} \hat{T}_R \\ \hat{B}_R \end{pmatrix}$$

$$\begin{pmatrix} \hat{U}_R \\ \hat{D}_R \end{pmatrix}$$

$$\begin{pmatrix} \hat{X}_R \\ \hat{Y}_R \end{pmatrix}$$

$$\begin{pmatrix} T_L \\ B_L \\ t_L \\ b_L \\ t'_R \end{pmatrix}_{\frac{2}{3}}$$

$$\begin{pmatrix} U_L \\ D_L \\ X_L \\ Y_L \\ b'_R \end{pmatrix}_{-\frac{1}{3}}$$

$$\Psi(x, -y) = P_0 \gamma^5 \Psi(x, y)$$

Brane scalar

$$\hat{\Phi} \left(0, \frac{1}{2}\right)$$

$$\left(\frac{1}{2}, 0\right)$$

vector rep

$$\left(\frac{1}{2}, \frac{1}{2}\right) \oplus (0, 0)$$

$$\begin{pmatrix} \hat{L}_{2XR} \\ \hat{L}_{2YR} \end{pmatrix}$$

$$\begin{pmatrix} \hat{L}_{3XR} \\ \hat{L}_{3YR} \end{pmatrix}$$

$$\begin{pmatrix} \hat{L}_{1XR} \\ \hat{L}_{1YR} \end{pmatrix}$$

$$\begin{pmatrix} \nu_{\tau L} \\ \tau_L \\ L_{1XL} \\ L_{1YL} \\ \tau'_R \end{pmatrix}_{-1}$$

$$\begin{pmatrix} L_{2XL} \\ L_{2YL} \\ L_{3XL} \\ L_{3YL} \\ \nu'_{\tau R} \end{pmatrix}_0$$

Leptons

Matter content

Planck brane

TeV brane

$$\begin{pmatrix} \hat{T}_R \\ \hat{B}_R \end{pmatrix}$$

$$\begin{pmatrix} \hat{U}_R \\ \hat{D}_R \end{pmatrix}$$

$$\begin{pmatrix} \hat{X}_R \\ \hat{Y}_R \end{pmatrix}$$

$$\left(\frac{1}{2}, 0\right)$$

$$\begin{pmatrix} \hat{L}_{2XR} \\ \hat{L}_{2YR} \end{pmatrix}$$

$$\begin{pmatrix} \hat{L}_{3XR} \\ \hat{L}_{3YR} \end{pmatrix}$$

$$\begin{pmatrix} \hat{L}_{1XR} \\ \hat{L}_{1YR} \end{pmatrix}$$

$$\begin{pmatrix} T_L \\ B_L \\ t_L \\ b_L \\ t'_R \end{pmatrix}_{\frac{2}{3}}$$

$$\begin{pmatrix} U_L \\ D_L \\ X_L \\ Y_L \\ b'_R \end{pmatrix}_{\frac{1}{3}}$$

$$\left(\frac{1}{2}, \frac{1}{2}\right) \oplus (0, 0)$$

$$\begin{pmatrix} \nu_{\tau L} \\ \tau_L \\ L_{1XL} \\ L_{1YL} \\ \tau'_R \end{pmatrix}_{-1}$$

$$\begin{pmatrix} L_{2XL} \\ L_{2YL} \\ L_{3XL} \\ L_{3YL} \\ \nu'_{\tau R} \end{pmatrix}_0$$

$$\langle \hat{\Phi} \rangle \neq 0$$

$$\left(0, \frac{1}{2}\right)$$

SM spectrum

Anomaly cancellation

## Effective interactions

$$\text{AB phase} \quad \hat{\theta}_H = \theta_H + \frac{H}{f_H} \quad f_H = \frac{2}{\sqrt{kL}} \frac{m_{KK}}{\pi g}$$

$$\mathcal{L}_{\text{eff}} \sim -V_{\text{eff}}(\hat{\theta}_H)$$

YH 1983, Oda-Weiler 2005

Falkowski 2007

$$-m_W (\hat{\theta}_H)^2 W_\mu^\dagger W^\mu - \frac{1}{2} m_Z (\hat{\theta}_H)^2 Z_\mu Z^\mu$$

Sakamura-YH 2006, 2007

$$-m_f (\hat{\theta}_H) \bar{\psi}_f \psi_f$$

YH-Kobayashi 2008

$$\theta_H \sim \theta_H + 2\pi$$

## Gauge-Higgs

$$m_W(\hat{\theta}_H) \sim \frac{1}{2} g \underline{f_H \sin \hat{\theta}_H}$$

$$m_Z(\hat{\theta}_H) \sim \frac{1}{2 \cos \theta_W} g \underline{f_H \sin \hat{\theta}_H}$$

$$m_f(\hat{\theta}_H) \sim y_f \underline{f_H \sin \hat{\theta}_H}$$

$$\hat{\theta}_H = \theta_H + \frac{H}{f_H}$$

*periodic  
nonlinear*

## SM

$$\frac{1}{2} g \underline{(v + H)}$$

$$\frac{1}{2 \cos \theta_W} g \underline{(v + H)}$$

$$y_f \underline{(v + H)}$$





WWH  
ZZH  
Yukawa

$$= SM \times \cos \theta_H$$

WWHH  
ZZHH

$$= SM \times \cos 2\theta_H$$

# Effective potential



EW symmetry breaking by Hosotani mechanism

$$m_H : 70 \text{ GeV} \sim 135 \text{ GeV}$$

$$\theta_H = \frac{\pi}{2}$$



WWH, ZZH, Yukawa = 0

*LEP2 bound is evaded.*

# H parity



$H : -$   
all other SM particles :  $+$

**Proof (1)**      YH, Ko, Tanaka, 0908.0212

**Proof (2)**      YH, Tanaka, Uekusa, 1010.6135

$$SO(5) : \quad SO(4) \simeq SU(2)_L \times SU(2)_R \quad SO(5)/SO(4)$$

$$\{ T^\alpha \} = \quad \{ T^{a_L}, T^{a_R}, T^{\hat{a}}, T^{\hat{4}} \}$$

Algebra is invariant under

$$\Rightarrow \{ T'^\alpha \} = \quad \{ T^{a_R}, T^{a_L}, T^{\hat{a}}, -T^{\hat{4}} \}$$

$$T'^\alpha = \Omega_H T^\alpha \Omega_H^{-1} \quad \Omega_H = \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & & & -1 & \\ & & & & 1 \end{pmatrix}$$

$$P_H : \quad \begin{aligned} SU(2)_L &\leftrightarrow SU(2)_R \\ T^{\hat{4}} &\rightarrow -T^{\hat{4}} \end{aligned}$$

 **Agashe, Contino, Da Rold, Pomarol 2006**

$$\text{At } \theta_H = \frac{\pi}{2}$$

bulk action      invariant under  $A_M \rightarrow \Omega_H A_M \Omega^{-1}$

  $P_H$ -inv

brane interactions       $P_H$  odd fields do not couple.



Theory is H parity invariant.

$$P_H = \begin{cases} + & W^{(n)}, Z^{(n)}, \gamma^{(n)}, gluon^{(n)}, q^{(n)}, \ell^{(n)}, \dots \\ - & H^{(n)}, W'^{(n)}, Z'^{(n)}, q'^{(n)}, \ell'^{(n)}, \dots \end{cases}$$


**Higgs field** : the lightest  $P_H$ -odd field.

$WWH, ZZH, Yukawa = 0$

Stable

# Stable Higgs = Dark Matter





WMAP data  $\rightarrow m_H \sim 70 \text{ GeV}$


$T_f \sim 3 \text{ GeV}$

$z_L$	$10^5$	$10^{10}$	$10^{15}$
$m_H$	72 GeV	108	135

$HH \rightarrow b\bar{b}$  34%  
 $\rightarrow WW$  61%

# How to see the Higgs bosons at LHC/ILC

## Production:



Stable Higgs boson

=

missing energy,  
missing momentum

Cheung, Song, 1004.2783

Alves, 1008.0016

$\nu, \bar{\nu}$  background

hard at LHC, possible at ILC

## Gauge couplings precision measurements

EW universality is slightly broken in 4D.

Forward-backward asymmetry in  $e^+e^- \rightarrow Z \rightarrow \ell\bar{\ell}, q\bar{q}$

Z-decay widths (branching ratios)

	No. data	SM	$z_L : 10^{15}$	$z_L : 10^{10}$	$z_L : 10^5$
$\sin^2 \theta_W$		0.2312	0.2309	0.2303	0.2284
$\chi^2(AFB)$	6	10.8335	6.3044	6.4345	7.1251
$\chi^2(Z \text{ decay})$	8	13.6264	16.5153	37.6908	184.468




 $z_L \geq 10^{15}$



# KK modes spectrum and production

YH, Tanaka, Uekusa, in preparation

## 1st KK modes



	$z_L : 10^{15}$	$z_L : 10^5$
$m_{\text{KK}}$	1466	836
$Z^{(1)}$	1130	659
$gluon^{(1)}$	1144	678
$u^{(1)}$	1361	1037
$t^{(1)}$	1121	634

in GeV

couplings/ $g_w$ $z_L = 10^{15}$		
	$Z$	$Z^{(1)}$
$u_L$	0.3485	-0.0400
$t_L$	0.3219	-0.2058
$d_L$	-0.4260	0.0488
$b_L$	-0.4265	-0.5581
$u_R$	-0.1562	-1.6557
$t_R$	-0.1835	-1.4668
$d_R$	0.0781	0.8279
$b_R$	0.0781	0.7227
$e_L$	-0.2710	0.0311
$\mu_L$	-0.2710	0.0310
$e_R$	0.2344	2.5161
$\mu_R$	0.2343	2.4201

Stronger for right-handed

KK  $Z^{(1)}$

$z_L$	$10^{15}$	$10^5$
$m$	1130	653
$\Gamma$	385	94

in GeV

at Tevatron

$$p \bar{p} \rightarrow Z' \rightarrow e^+ e^-$$

signal

2 pb	for $z_L = 10^5$
0.3 pb	$10^{10}$
0.08 pb	$10^{15}$

Consistent for  $z_L \geq 10^{15}$

# KK gluon<sup>(1)</sup>

## Strong couplings for right-handed quarks

Couplings/ $g_s$

$u_R$	6.32
$c_R$	6.04
$t_R$	5.60

In naive perturbation theory  $\Gamma \sim 13 m$

Decays into  
light and heavy, right-handed quarks.

# Summary

$SO(5) \times U(1)$  gauge-Higgs unification

Higgs naturally becomes stable.

Dark Matter = Higgs  $\Rightarrow m_H \sim 70 \text{ GeV} \Rightarrow z_L \sim 10^5$

EW precision data (gauge couplings)


$$\Rightarrow z_L \geq 10^{15}$$

KK modes at Tevatron/LHC

$$\Rightarrow z_L \geq 10^{15}$$

# Direct detection rate

CDMS II



GHU  
stable Higgs

*many  
uncertainties*

Higgs = Dark Matter

Annihilation

